

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) **EP 0 398 692 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
21.08.1996 Bulletin 1996/34

(21) Application number: 90305303.1

(22) Date of filing: 16.05.1990

(51) Int. Cl.⁶: **C07C 251/48**, A01N 37/50,
C07C 323/57, C07C 317/44,
C07D 239/34, C07C 323/63,
C07F 7/10, C07D 213/643,
C07D 309/12, C07D 277/68,
C07D 215/227

(54) **Alkoxyiminoacetamide derivatives and their use as fungicides**

Alkoxyiminoacetamid-Derivate und ihre Verwendung als pilztötendes Mittel

Dérivés d'alkoxyiminoacétamide et leur application comme fongicides

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(30) Priority: 17.05.1989 JP 124059/89
29.12.1989 JP 341175/89

(43) Date of publication of application:
22.11.1990 Bulletin 1990/47

(60) Divisional application: 94111766.5

(73) Proprietor: SHIONOGI SEIYAKU KABUSHIKI
KAISHA
trading under the name of
SHIONOGI & CO. LTD.
Osaka 541 (JP)

(72) Inventors:
• Hayase, Yoshio
Kameyama-shi, Mie-ken (JP)
• Kataoka, Takahiro
7, Clayton, MO 63105 (US)
• Takenaka, Hideyuki
Koga-gun, Shiga-ken (JP)

• Ichinari, Mitsuhiro
Koga-gun, Shiga-ken (JP)
• Masuko, Michio
Koga-gun, Shiga-ken (JP)
• Takahashi, Toshio
Nishinomiya-shi, Hyogo-ken (JP)
• Tanimoto, Norihiko
Koga-gun, Shiga-ken (JP)

(74) Representative: Hardisty, David Robert et al
BOULT, WADE & TENNANT
27 Fumival Street
London EC4A 1PQ (GB)

(56) References cited:
EP-A- 0 023 891 EP-A- 0 253 213
EP-A- 0 254 426 EP-A- 0 299 382
US-A- 3 219 694

Remarks:

The file contains technical information submitted
after the application was filed and not included in
this specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description


The present invention relates to alkoxyiminoacetamide derivatives and their use as fungicides, particularly in the agricultural field.

Various alkoxyiminoacetic acid derivatives having a diaryl ether group are known to exert biological activities including fungicidal activity, herbicidal activity, etc. For instance, some phenoxyphenylalkoxyiminoacetic esters are disclosed to exert fungicidal activity in JP-A-63-23852 and JP-A-63-30463, some phenoxyphenylalkoxyiminoacetamides are disclosed to have herbicidal activity in JP-A-55-35006 and JP-A-56-29560, and some pyridyloxyphenyl-alkoxyiminoacetamides are disclosed to show herbicidal activity in JP-A-56-55368. However, the relationship between a chemical structure and a biological activity has not sufficiently been clarified so that it is still hardly possible to predict the biological activity to be exerted by a chemical compound from its chemical structure.


Furthermore, EP-A-23 891 discloses herbicidal compounds which overlap with the fungicidal compounds of the present invention, namely compounds wherein Z is O, and A is 2-pyridyl.

Also, EP-A-253 213 discloses a group of esters which may be considered to be related to the compounds of this invention, said prior compounds also having fungicidal activity.

As the result of an extensive study on alkoxyiminoacetic acid derivatives having a diaryl ether group, it has been found that a group of alkoxyiminoacetamides of the following formula show a strong fungicidal activity against a wide variety of fungi, particularly phyto-pathogenic fungi:



wherein R¹ and R² are each hydrogen, lower alkyl or cyclo(lower)alkyl; R³ is lower alkyl or cyclo(lower)alkyl; R⁴ and R⁵ are each hydrogen, lower alkyl, lower alkoxy, halogen-substituted lower alkyl, lower alkyl-substituted silyl, halogen or nitro; A represents an unsaturated hydrocarbon group, a halogen-substituted unsaturated hydrocarbon group, a phenyl group or a heterocyclic group, said phenyl group or heterocyclic group being optionally substituted with not more than three substituents; and Z is -CH₂-, -CH(OH)-, -CO-, -O-, -S-, -SO-, -NR- (R being hydrogen or lower alkyl), -CH₂CH₂-, -CH=CH-,



-CH₂O-, -CH₂S-, -CH₂SO-, -OCH₂-, -SCH₂- or -SOCH₂-, provided that when Z is O A is other than 2-pyridyl

In this specification, the term "lower" is used to mean a group having not more than 8 carbon atoms, preferably not more than 6 carbon atoms. Preferably, the term "lower alkyl" means an alkyl group having not more than 6 carbon atoms, more preferably not more than 4 carbon atoms (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl). The term "cyclo(lower)alkyl" means a cycloalkyl group having 3 to 8 carbon atoms, preferably 3 to 6 carbon atoms (e.g. cyclopropyl, cyclopentyl, cyclohexyl). Preferably the term "lower alkoxy" means an alkoxy group having not more than 6 carbon atoms, more preferably not more than 4 carbon atoms (e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy). The term "lower alkenyl" refers to an alkenyl group having 2 to 8 carbon atoms, preferably 3 to 6 carbon atoms (e.g. allyl, isopropenyl, butenyl, isobutenyl, pentenyl, hexenyl, hexadienyl).


The term "halogen" covers chlorine, bromine, iodine and fluorine. The terms "halogen-substituted", "lower alkyl-substituted" represents any group substituted with not more than three halogen atoms or lower alkyl groups.

The unsaturated hydrocarbon group represented by the symbol A means an alkenyl or alkadienyl group having 2 to 12 carbon atoms, preferably 3 to 10 carbon atoms. The heterocyclic group represented by the symbol A may be a 5- to 7-membered, heteromonocyclic group having not more than three ring nitrogen atoms, and specific examples are


pyridine, pyrimidine, pyridazine. When said phenyl or heterocyclic group is substituted, the substituent(s) may be one to three chosen from lower alkyl, lower alkanoyl, lower alkyl-substituted silyl, halogen-substituted alkyl, di(lower)alkylamino, phenyl, phenyl(lower)alkyl, phenyl(lower)alkenyl, furyl(lower)alkyl, furyl(lower)alkenyl, halogen, nitro, cyano, -OR⁶ (in which R⁶ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower alkanoyl, phenyl, lower alkoxyphenyl, nitrophenyl, phenyl(lower)alkyl, cyanophenyl(lower)alkyl, benzoyl, tetrahydropyranyl, pyridyl, trifluoromethylpyridyl, pyrimidyl, benzothiazolyl, quinolyl, benzoyl(lower)alkyl, benzenesulfonyl or lower alkylbenzenesulfonyl) and -CH₂-Z'-R⁷ (in which Z' is -O-, -S-, -SO- or -NR- (R being hydrogen or lower alkyl) and R⁷ is phenyl, halophenyl, lower alkoxyphenyl, pyridyl or pyrimidinyl).

A main object of the present invention is to provide the alkoxyiminoacetamides of the formula (I) as novel chemical substances. Another object of this invention is to provide the alkoxyiminoacetamides (I) useful as fungicides, particularly as an agricultural fungicide. A further object of the invention is provide a fungicidal composition comprising at least one of the alkoxyiminoacetamides (I) as the active ingredient. A still further object of the invention is to provide a process for preparing the alkoxyiminoacetamides (I). These and other objects of the invention will be apparent to those skilled in the art from the foregoing and subsequent descriptions.


Within the formula (I), there are included the alkoxyiminoacetamides of the following formulas:




(I-1)






(I-2)



(I-3)




(I-4)




wherein R^1 and R^2 are each hydrogen or lower alkyl (particularly methyl), R^3 is lower alkyl (particularly methyl), R^4 and R^5 are each hydrogen, lower alkyl (particularly methyl), lower alkoxy (particularly methoxy), halogen-substituted lower alkyl (particularly trifluoromethyl), lower alkyl-substituted silyl (particularly trimethylsilyl), halogen (particularly, chlorine, iodine or fluorine) or nitro and X_1 , X_2 and X_3 are each hydrogen, lower alkyl (particularly methyl, ethyl, propyl, isopropyl, butyl or t-butyl), lower alkoxy (particularly methoxy), lower alkanoyl (particularly acetyl), halogen-substituted lower alkyl (particularly trifluoromethyl), lower alkyl-substituted silyl (particularly trimethylsilyl), halogen (particularly chlorine, bromine or fluorine), nitro, cyano, di(lower)alkylamino (particularly dimethylamino), phenyl, phenyl(lower)alkenyl, (particularly phenylethenyl), furyl(lower)alkenyl (particularly furylethenyl), hydroxyl, lower alkynyloxy (particularly 2-propynyloxy), lower alkanoyloxy (particularly acetoxy), benzyloxy, phenoxy, lower alkoxyphenoxy (particularly methoxyphenoxy), nitrophenoxy, benzyloxy, cyanobenzyloxy, tetrahydropyranyloxy, pyrimidinyl, pyridyl, trifluoromethylpyridyl, benzothiazolyl, quinolyl, benzoyl(lower)alkoxy (particularly benzoylmethoxy), benzenesulfonyloxy or lower alkylbenzenesulfonyloxy (particularly toluenesulfonyloxy).

In addition to the above alkoxyiminoacetamides (I), there are also included the following ones:




wherein R^1 and R^2 are each hydrogen or lower alkyl (particularly methyl), R^3 is lower alkyl (particularly methyl), R^4 and R^5 are each hydrogen, lower alkyl (particularly methyl), lower alkoxy (particularly methoxy), halogen-substituted lower alkyl (particularly trifluoromethyl), lower alkylsubstituted silyl (particularly trimethylsilyl), halogen (particularly, chlorine, iodine or fluorine) or nitro, R^7 is phenyl, lower alkoxyphenyl (particularly methoxyphenyl), halophenyl (particularly bromophenyl), pyridyl or pyrimidinyl and Z' is $-O-$, $-S-$, $-SO-$, or $-NR-$ (in which R is lower alkyl, particularly methyl);




wherein R^1 and R^2 are each hydrogen or lower alkyl (particularly methyl), R^3 is lower alkyl (particularly methyl),




A is pyridyl or pyrimidinyl, Z is $-O-$, $-OCH_2-$ or $-CH_2O-$ and X_1 is hydrogen or trifluoromethyl;



wherein R^1 and R^2 are each hydrogen or lower alkyl (particularly methyl), R^3 is lower alkyl (particularly methyl) and Z is $-CH_2CH_2-$,




-CH=CH-, -CH(OH)- or -CO-; and




wherein R^1 and R^2 are each hydrogen or lower alkyl (particularly methyl), R^3 is lower alkyl (particularly methyl) and A is alkenyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms (particularly chlorine) or alkadienyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms (particularly chlorine).

The alkoxyiminoacetamides (I) may be produced by various processes, among which a typical standard process starts from the corresponding oxocarboxylic acid of the formula:



wherein A and Z are each as defined above and substantially comprises amidation and alkoxylation (alkoxime-formation), or vice versa. Namely, the core portion of such process is represented by Scheme A:

Scheme A



wherein R^1 , R^2 , R^3 , R^4 , R^5 , A and Z are each as defined above.

Explaining the conversions in Scheme A, the oxocarboxylic acid (II) or its derivative on the carboxyl group is reacted with an amine of the formula:




wherein R^1 and R^2 are each as defined above (amidation), and the resultant oxocarbonamide (III) is reacted with an alkoxyamine of the formula:




wherein R^3 is as defined above (alkoximation) to give the objective alkoxyiminoacetamide (I).

When R^1 in the amine (a) is hydrogen, the reaction proceeds not only on the carboxyl group but also on the carbonyl group in the oxocarboxylic acid (II) or its derivative on the carboxyl group so that the following iminocarbonamide is by-produced:



wherein R^2 , R^4 , R^5 , A and Z are each as defined above. Particularly when such amine (a) is used in two moles or more to one mole of the oxocarboxylic acid (II) or its derivative, the iminocarbonamide (III') is obtainable as a main product. Not only the oxocarbonamide (III) but also the iminocarbonamide (III') can be converted into the objective alkoxyiminoacetamide (I), when reacted with the alkoxyamine (b).

When desired, the alkoximation may be effected in two steps, i.e. reaction of the oxocarbonamide (III) or the iminocarbonamide (III') with hydroxylamine and reaction of the resultant hydroxyiminoacetamide (IV) with an alkylating agent of the formula:



wherein X is an acid residue such as a halogen atom (e.g. chlorine, bromine) or a sulfonyl group (e.g. methanesulfonyloxy, ethanesulfonyloxy, toluenesulfonyloxy).

Alternatively, the oxocarboxylic acid (II) or its derivative on the carboxyl group is reacted with the alkoxyamine (b) (alkoximation), and then the resulting alkoxyiminocarboxylic acid (V) or its derivative on the carboxyl group is reacted with the amine (a) (amidation) to give the objective alkoxyiminoacetamide (I).

When desired, the alkoximation may be effected in two steps, i.e. reaction of the oxocarboxylic acid (II) or its derivative on the carboxyl group with hydroxylamine and reaction of the resulting hydroxyiminocarboxylic acid (VI) or its derivative on the carboxyl group with the alkylating agent (c).

The thus produced alkoxyiminocarboxylic acid (V) or its derivative on the carboxyl group is then reacted with the amine (a) to give the objective alkoxyiminoacetamide (I).

As understood from the above, the oxocarboxylic acid (II) as the starting material in the above process may be used as the free acid or its derivative. Examples of the derivative are acid esters, acid anhydrides, acid halides, etc.

Any reaction included in the above process such as the amidation (reaction with the amine (a)), the alkoximation (reaction with the alkoxyamine (b)), the oximation (reaction with hydroxylamine) and the alkylation (reaction with the alkylating agent (c)) is per se conventional and may be carried out in a manner as commonly known to those skilled in the art. In general, those reactions are performed in a solvent, preferably under conditions to facilitate or promote their proceeding at a temperature of room temperature to the reflux temperature of the reaction mixture, e.g. from 10 to 200°C.

The condition to facilitate or promote the proceeding of the reaction may be appropriately taking into consideration the kinds of the reactants or the kinds of the by-products to be eliminated from the reactants. For instance, the amidation proceeds between a carboxylic acid or its derivative on the carboxyl group such as an acid ester (e.g. methyl ester, ethyl ester, cyanomethyl ester, p-nitrophenyl ester), an acid anhydride (e.g. a mixed acid anhydride with trichloroacetic acid) or an acid halide (e.g. acid chloride, acid bromide) and the amine (a) to give a carbonamide together with an alco-


hol, an acid or the like. In order to eliminate these by-products, a condensing agent (e.g. 1,3-dicyclohexylcarbodiimide, N-cyclohexyl-N'-morpholinoethylcarbodiimide, 1,3-diisopropylcarbodiimide, methyl chlorocarbonate, ethyl chlorocarbonate), a dehydrating agent (e.g. thionyl chloride, sulfuryl chloride, phosphorus oxychloride, phosphorus tribromide, phosphorus pentachloride, polyphosphoric acid), an acid-eliminating agent (e.g. pyridine, triethylamine, sodium methoxide, sodium hydroxide, sodium carbonate), etc. may be used. Further, for instance, the alkoxylation proceeds between a ketone and the alkoxyamine (b) to give an alkoxime together with water. Usually, a base such as sodium hydroxide, sodium acetate or pyridine is incorporated into the reaction system.

The reaction is normally effected in a solvent appropriately chosen depending upon the kind of the reaction from dioxane, methylene chloride, chloroform, diethyl ether, tetrahydrofuran, acetone, dimethylformamide, dimethylsulfoxide, pyridine, acetonitrile, benzene, toluene, xylene, etc.


As stated above, Scheme A shows only the core portion of the typical standard process, and it is of course possible to produce the oxocarbonamide (III) or the alkoxyiminocarboxylic acid (V) by any other process not going through the oxocarboxylic acid (II) in a conventional manner. Also, the once produced alkoxyiminoacetamide (I) may be subjected to conversion on any substituent on the benzene ring such as represented by R⁴, R⁵ or -Z-A.

Referring to Scheme 1 to 11 as hereinafter given, some specific procedures for production of the alkoxyiminoacetamides (I) will be explained in details. In Scheme 1 to 11, Me is methyl, Hal is halogen, X is an acid residue such as halogen, Z' is -O-, -S- or -NR- (R being lower alkyl) and R⁴, R⁵, R⁶, R⁷, X₁, X₂ and X₃ are each as defined above.


Scheme 1:




Scheme 2:




Scheme 3:




Scheme 4:




Scheme 5:




Scheme 6:




Scheme 7:




Scheme 8:




Scheme 9:



Scheme 10:



Scheme 11:



In Scheme 1, the starting diphenyl ether is reacted with butyl lithium to introduce lithium into the 2 position on the benzene ring, followed by reaction with dimethyl oxalate to give a ketonic ester as an example of the derivative of the oxocarboxylic acid (II). Then, the ketonic ester is reacted with methoxyamine to give a methoxyiminoacetate acid ester, which is an example of the derivative of the alkoxyiminocarboxylic acid (V).

In Scheme 2, the starting phenol is reacted with a nitrophenyl halide in the presence of a base (e.g. sodium amide, potassium carbonate) in an inert solvent (e.g. dimethylsulfoxide, dimethylformamide) to give a diphenyl ether. The diphenyl ether is subjected to reduction on the nitro group, and the resultant amine is subjected to Sandmeyer's reaction for conversion of the amino group into a halogen atom. Then, the halogenated compound is reacted with butyl lithium, followed by condensation with dimethyl oxalate to a ketonic ester. This ketonic ester is an example of the derivative of the oxocarboxylic acid (II).

In Scheme 3, the starting phenol is reacted with a benzyl halide in the presence of a base (e.g. sodium amide, triethylamine, sodium hydroxide, barium oxide, silver oxide, sodium hydride) in an inert solvent (e.g. ether, dimethylsulfoxide, dimethylformamide, tetrahydrofuran) to give a benzyl phenyl ether, which is an example of the derivative of the alkoxyiminocarboxylic acid (V).

In Scheme 4, the starting benzyl halide, obtained from the corresponding tolyl compound by the reaction with a halogenating agent (e.g. chlorine, bromine, t-butyl hypohalogenite, N-halosuccinimide, trichloromethanesulfonyl halide) at a high temperature or in the presence of light or a peroxide, is reacted with a phenol in the same manner as in Scheme 3 to give a phenyl benzyl ether as an example of the derivative of the alkoxyiminocarboxylic acid (V).

In Scheme 5, methyl 2-tolylacetate is oxidized with selenium oxide, or 2-tolyl halide is reacted with butyl lithium or magnesium, followed by reaction with dimethyl oxalate, whereby a ketonic ester is obtained. The ketonic ester is reacted with methoxyamine hydrochloride to give a methoxime. Alternatively, the ketonic ester is first reacted with hydroxylamine hydrochloride, and then the resultant oxime is reacted with dimethyl sulfate in the presence of a base to give a methoxime. The thus produced methoxime is subjected to halogenation so that the methyl group on the benzene ring is converted into a halomethyl group. The resultant benzyl halide is then reacted with a reagent of the formula: A-Z"-H (wherein Z" is -O-, -S- or -NR- (R being lower alkyl)) in a reactive form in the presence of a base to give a methoxyiminoacetic acid ester, which is an example of the derivative of the alkoxyiminocarboxylic acid (V).

In Scheme 6, the starting tetrahydropyranyloxyphenyl halide is reacted with butyl lithium, followed by reaction with dimethyl oxalate to give a ketonic ester. The ketonic ester is then reacted with methoxyamine hydrochloride or with hydroxylamine hydrochloride and dimethyl sulfate in order to give a tetrahydropyranyl phenyl ether, which is an example of the derivative of the alkoxyiminocarboxylic acid (V). The tetrahydropyranyl phenyl ether is then treated with methylamine, whereby the carboxylic ester is converted into a carbonamide simultaneously with elimination of the tetrahydropyranyl group. The resultant phenol is reacted with an alkenyl or alkadienyl halide in the presence of a base to give an alkenyl or alkadienyl phenyl ether, which is an example of the alkoxyiminoacetamide (I).

In Scheme 7, the starting hydroxyphenyl halide is reacted with an aryl halide (e.g. phenyl bromide, pyridyl bromide) to give an aryloxyphenyl halide. The aryloxyphenyl halide is converted into an aryl phenyl ether in the same manner as in Scheme 6, i.e. by reaction with butyl lithium or magnesium, followed by treatment with dimethyl oxalate and reaction with methoxyamine hydrochloride or with hydroxylamine hydrochloride and then dimethyl sulfate. The aryl phenyl ether is an example of the derivative of the alkoxyiminocarboxylic acid (V).

In Scheme 8, the starting iminoacetic acid ester is reacted with methylamine and an alkali metal acetate, whereby the ester group is converted into a carbonamide group and the halomethyl group is converted into a hydroxymethyl group. The resultant hydroxyphenylcarbonamide is reacted with an aryl halide (e.g. phenyl halide, pyrimidinyl halide) in the presence of a base to give an aryl benzyl ether, which is an example of the alkoxyiminoacetamide (I).

In Scheme 9, the starting dihydroxybenzene is reacted with 2-halonitrobenzene in the presence of a base, and the resultant nitrodiphenyl ether is subjected to reduction and Sandmeyer reaction in order to give 2-(hydroxyphenoxy)phenyl halide, which is then reacted with dihydropyran in the presence of an acid catalyst (e.g. sulfuric acid, hydrochloric acid, boron trifluoride). The resultant tetrahydropyranyloxyphenyloxyphenyl halide is reacted with butyl lithium or magnesium and then with dimethyl oxalate to give a ketonic ester as an example of the derivative of the oxocarboxylic acid (II). This ketonic ester is reacted with methoxyamine hydrochloride to give a methoxime as an example of the derivative of the alkoxyiminocarboxylic acid (V). Thereafter, the methoxime is reacted with R⁶X (wherein R⁶ and X are each as defined above) (e.g. phenyl halide, pyridyl halide, pyrimidinyl halide, quinolyl halide, benzothiazolyl halide, phenacyl halide, tosyl halide, acetyl halide, benzoyl halide, benzyl halide, propargyl halide, styryl halide) so as to convert the hydroxyl group into an R⁶O- group.

In Scheme 10, the starting methoxime is halogenated to give the corresponding halomethyl compound, which is then treated with an alkali metal acetate and methylamine to give a methoxyiminocarbonamide as an example of the alkoxyiminoacetamide (I). This methoxyiminocarbonamide is reacted with a group-introducing reagent of the formula: R⁷X in the presence of a base, whereby the hydroxyl group is converted into -OR⁷. Alternatively, said halomethyl compound is reacted with a reagent of the formula: R⁷-Z'-H (wherein R⁷ and Z' are each as defined above) and methylamine to give a methoxyiminocarbonamide as another example of the alkoxyiminoacetamide (I).

In Scheme 11, the starting phenyl halide is reacted with butyl lithium or magnesium and then with dimethyl oxalate to give a ketonic ester as an example of the derivative of the oxocarboxylic acid (II). The ketonic ester is reacted with methoxyamine hydrochloride or with hydroxylamine and dimethyl sulfate in order to give a methoxime as an example of the derivative of the alkoxyiminocarboxylic acid (V).

The alkoxyiminoacetamides (I) thus produced are usually obtained as a mixture of the E and Z forms, which can be separated into each of those forms.

The alkoxyiminoacetamides (I) show a strong fungicidal activity against a wide variety of phytopathogenic fungi on crop plants (e.g. rice plant, wheat, barley, rye, corn, common millet, millet, buck wheat, soybean, redbean, peanut), vegetables (e.g. cucumber, eggplant, tomato, pumpkin, kidney bean), fruit trees (e.g. citrus fruits, grape, apple, pear, peach), etc. They also show a fungicidal activity against phytopathogenic fungi in soil. Examples of the phytopathogenic fungi on which the alkoxyiminoacetamides (I) exert their fungicidal activity are *Pyricularia oryzae*, *Rhizoctonia solani*,

Erysiphe graminis, Sphaerotheca fuliginea, Erysiphe cichoracearum, Phytophthora infestans, Pseudoperonospora cubensis, Peronospora manshurica, Plasmopara viticola, Botrytis cinerea, Pythium aphanidermatum, Sclerotinia sclerotiorum, Corticium rolfsii, etc. Therefore, the alkoxyiminoacetamides (I) are useful as agricultural fungicides.

Application of the alkoxyiminoacetamides (I) may be performed before and/or after the infection with phytopathogenic fungi on plants. Application may be made to plants by any conventional procedure such as atomizing, scattering or spreading. Application may be also made through treatment of seeds of plants, soil where plants grow, paddy field for seedling or water for perfusion with the alkoxyiminoacetamides (I).

For practical usage, the alkoxyiminoacetamides (I) may be applied as such or in formulations such as solutions, emulsions, dispersions, dusts, wettable powders, oil sprays, emulsifiable concentrates, tablets, granules, fine granules, aerosols or flowables. Such formulations can be prepared in a conventional manner by mixing at least one of the alkoxyiminoacetamides (I) with an appropriate solid or liquid carrier(s) and, if necessary, an appropriate adjuvant(s) (e.g. surfactants, adherents, dispersants, stabilizers) for improving the dispersibility and other properties of the active ingredient.

Examples of solid carriers or diluents are botanical materials (e.g. flour, tobacco stalk powder, soybean powder, walnut-shell powder, vegetable powder, saw dust, bran, bark powder, cellulose powder, vegetable extract residue), fibrous materials (e.g. paper, corrugated cardboard, old rags), synthetic plastic powders, clays (e.g. kaolin, bentonite, fuller's earth), talcs, other inorganic materials (e.g. pyrophyllite, sericite, pumice, sulfur powder, active carbon), chemical fertilizers (e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, ammonium chloride), etc.

Examples of liquid carriers or diluents are water, alcohols (e.g. methanol, ethanol), ketones (e.g. acetone, methyl-ethylketone), ethers (e.g. diethyl ether, dioxane, cellosolve, tetrahydrofuran), aromatic hydrocarbons (e.g. benzene, toluene, xylene, methyl-naphthalene), aliphatic hydrocarbons (e.g. gasoline, kerosene, lamp oil), esters, nitriles, acid amides (e.g. dimethylformamide, dimethylacetamide), halogenated hydrocarbons (e.g. dichlorocarbon tetrachloride), etc.

Examples of surfactants are alkyl sulfuric esters, alkyl sulfonates, alkylaryl sulfonates, polyethylene glycol ethers, polyhydric alcohol esters, etc. Examples of adherents or dispersants may include casein, gelatin, starch powder, carboxymethyl cellulose, gum arabic, alginic acid, lignin, bentonite, molasses, polyvinyl alcohol, pine oil and agar. As stabilizers, there may be used PAP (isopropyl acid phosphate mixture), tricresyl phosphate (TCP), tolu oil, epoxydized oil, various surfactants, various fatty acids and their esters, etc.

In addition to the above components, other fungicides, insecticides, herbicides, fertilizers, etc. may be incorporated into the composition.

The composition as above formulated generally contains at least one of the alkoxyiminoacetamides (I) in a concentration of about 1 to 95 % by weight, preferably of about 2.0 to 80 % by weight. By using such composition as such or in a diluted form, the alkoxyiminoacetamides (I) are generally applied in such amounts as about 1.0 g to 5 kg/hectare, preferably about 2 to 100 g/hectare, usually in concentrations ranging from about 1 to 50,000 ppm, preferably from about 100 to 5,000 ppm.

Practical and presently preferred embodiments of the invention are illustratively shown in the following examples wherein the abbreviations indicate the following meanings: Me, methyl; OMe, methoxy; SiMe₃, trimethylsilyl; Me₂N, dimethylamino; (i)Pr, isopropyl; (i)PrO, isopropoxy; Ph, phenyl; OPh, phenoxy; tBu, t-butyl.

Example 1

Production of 2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 1):-

To a solution of methyl-(2-phenoxyphenyl)-2-oxoacetate (0.70 g) in methanol (10 ml), 28 % aqueous ammonium hydroxide (2.0 ml) was added, and the resultant mixture was stirred overnight, followed by removal of the solvent under reduced pressure. The residue was combined with diethyl ether (150 ml) and water (50 ml) under stirring and, upon dissolution, allowed to stand. The organic layer was separated, washed and dried over anhydrous sodium sulfate. Upon filtration, the filtrate was concentrated under reduced pressure to give residual yellow crystals, which were combined with methanol (10 ml) and O-methylhydroxylamine hydrochloride (1.49 g) while stirring and refluxed for 6 hours. Water (50 ml) was added to the reaction mixture, which was extracted with diethyl ether. The organic layer was washed with water and dried over anhydrous sodium sulfate, followed by filtration. The filtrate was concentrated under reduced pressure to give an oily residue. The residue was purified by silica gel column chromatography with benzene to give 0.36 g of the objective compound. $n_D^{23.3}$ 1.5978.

Elementary analysis (%) for C₁₅H₁₄H₂O₃.1/2H₂O:

Calcd.:	C, 64.50;	H, 5.41;	N, 10.03.
Found:	C, 64.63;	H, 5.13;	N, 9.89.

Example 2

- 5 Production of N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (isomeric mixture of Compound No. 3 and Compound No. 4):-

To a solution of nethyl-2-phenoxyphenyl-2-oxoacetate (0.70 g) in methanol (10 ml), 5 % aqueous methylamine (3.4 ml) was added, and the resultant mixture was stirred overnight, followed by removal of the solvent under reduced pressure. The residue was combined with diethyl ether (150 ml) and water (50 ml) under stirring and, upon dissolution, allowed to stand. The organic layer was separated, washed and dried over anhydrous sodium sulfate. Upon filtration, the filtrate was concentrated under reduced pressure to give an oily residue, which was purified by silica gel column chromatography with a mixture of benzene and ethyl acetate (100 : 5). High polar eluate was concentrated under reduced pressure to give an oily residue (0.55 g), which was combined with methanol (10 ml) and O-methyl hydroxylamine hydrochloride (0.40 g) under stirring and refluxed for 7 hours. Water (50 ml) was added to the reaction mixture, which was extracted with diethyl ether (150 ml). The extract was washed with water, dried and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with a mixture of benzene and ethyl acetate (100 : 5), whereby 0.25 g of Compound No. 3 was obtained as white crystals (m.p., 63 - 64.5°C) from the first eluate and 0.15 g of Compound No. 4 was obtained as white crystals (m.p., 48 - 51°C) from the second eluate.

Elementary analysis (%) for $C_{16}H_{16}N_2O_3 \cdot 1/5H_2O$:			
Compound No. 3			
Calcd.:	C, 66.74;	H, 5.74;	N, 9.73.
Found:	C, 66.75;	H, 5.68;	N, 9.71.
Compound No. 4			
Calcd.:	C, 66.74;	H, 5.74;	N, 9.73.
Found:	C, 66.85;	H, 5.90;	N, 9.66.

Example 3

Production of N,N-dimethyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 2):-

To a solution of methyl-2-(2-phenoxyphenyl)-2-oxoacetic acid (0.70 g) in methanol (10 ml), 50 % aqueous dimethylamine (2.5 ml) was added, and the resultant mixture was stirred overnight. The reaction mixture was treated in the same manner as in Example 1 to give an oily residue (0.50 g). The residue was combined with methanol (10 ml) and O-methylhydroxylamine hydrochloride (0.34 g), followed by stirring for 7 hours. The reaction mixture was treated in the same manner as in Example 2 to give 0.26 g of the objective compound as white crystals. m.p., 63 - 64°C.

Elementary analysis (%) for $C_{17}H_{18}N_2O_3$:			
Calcd.:	C, 68.44;	H, 6.08;	N, 9.38.
Found:	C, 68.44;	H, 6.21;	N, 9.48.

Example 4

Production of N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 3):-

A solution of N-methyl-2-(2-phenoxyphenyl)-2-oxoacetamide (1.50 g) and hydroxylamine hydrochloride (0.69 g) in

methanol (10 ml) was refluxed for 220 minutes while stirring, and water (75 ml) was added to the reaction mixture, followed by extraction with dichloromethane twice (150 ml and 100 ml). The organic layer was washed with water (75 ml) and dried over anhydrous sodium sulfate. Upon filtration, the filtrate was concentrated under reduced pressure to give a crystalline residue. The residue was combined with dimethylformamide (DMF) (7 ml), potassium carbonate (1.6 g) and dimethyl sulfate (0.58 ml) and stirred overnight. The reaction mixture was treated in the same manner as in Example 2 to give 0.80 g of the objective compound as an isomeric product.

Example 5

10 Production of (Z)-N-methyl-2-[2-(3-tolyloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 5):-

(1) To a solution of 3-phenoxytoluene (2.00 g) in diethyl ether (30 ml), a solution of butyl lithium in hexane (14.1 mmol) was added at 0°C, followed by stirring at room temperature for 12 hours. A solution of dimethyl oxalate (2.57 g) in diethyl ether (30 ml) was dropwise added thereto at 0°C, and the resultant mixture was stirred for 5 hours and neutralized with 1N hydrochloric acid, followed by extraction with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give a mixture of the following products, of which ¹H-NMR was measured in chloroform-d₁ (CDCl₃) with 270 MHz spectrometer and chemical shifts are reported in ppm down field from tetramethylsilane as an internal standard. The coupling constant (J) is to be represented by Hz. Abbreviations specifically denotes the following meanings: s, singlet; d, doublet; t, triplet; q, quartet; sept, septet; brs, broad singlet; brd, broad doublet; m, multiplet.

Product (A):-

Methyl-2-[2-(3-tolyloxy)phenyl]-2-oxoacetate (yield, 965 mg)

¹H-NMR: 2.35 (3H, s), 3.72 (3H, s), 6.86 (1H, d, J = 7.8), 6.88 (1H, s), 7.00 (1H, d, J = 7.8), 7.17 - 7.28 (3H, m), 7.51 (1H, td, J = 7.8, 2.0), 7.96 (1H, dd, J = 7.8, 2.0);

Product (B):-

Methyl-2-(4-methyl-2-phenoxyphenyl)-2-(4-methyl-2-phenoxyphenyl)-2-oxoacetate (yield, 260 mg)

¹H-NMR: 2.32 (3H, s), 3.69 (3H, s), 6.65 (1H, s), 7.03 - 7.11 (2H, m), 7.14 - 7.19 (2H, m), 7.34 - 7.38 (2H, m), 7.88 (1H, d, J = 7.8).

(2) Product (A) (965 mg), i.e. methyl-2-[2-(3-tolyloxy)phenyl]-2-oxoacetate as obtained (1) above, and O-methylhydroxylamine hydrochloride (597 mg) were dissolved in methanol (2 ml), and the resultant mixture was heated under reflux for 5 hours. The reaction mixture was cooled to room temperature, followed by removal of ethanol. The residue was added to water and extracted with diethyl ether. The solvent was dried and evaporated to give a mixture of the following products:

Product (A'):-

Methyl-(Z)-2-[2-(3-tolyloxy)phenyl]-2-methoxyiminoacetate (yield, 286 mg)

¹H-NMR: 2.32 (3H, s), 3.65 (3H, s), 4.02 (3H, s), 6.75 - 6.94 (3H, m), 7.12 - 7.36 (4H, m), 7.83 (1H, d, J = 7.8);

Product (B'):-

Methyl-(E)-2-(3-tolyloxy)phenyl]-2-methoxyiminoacetate (739 mg)

¹H-NMR: 2.32 (3H, s), 3.78 (3H, s), 4.03 (3H, s), 6.77 - 6.92 (3H, m), 7.11 - 7.38 (5H, m).

(3) To Product (A') (286 mg), i.e. methyl-(Z)-2-[2-(3-tolyloxy)phenyl]-2-methoxyiminoacetate, 30 % methanolic methylamine (197 mg) was added, followed by stirring at room temperature for 12 hours. Excess amine and methanol were removed from the mixture, which was subjected to silica gel column chromatography with a mixture of hexane and ethyl acetate to give 201 mg of (Z)-N-methyl-2-[2-(3-tolyloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 5).

Example 6

Production of (E)-N-methyl-2-[2-(3-tolylloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 6):-

- 5 In the same manner as in Example 5, the objective compound was prepared from methyl-(E)-2-[2-(3-tolylloxy)phenyl]-2-methoxyiminoacetate (Product (B)).

Example 7

- 10 Production of (Z)-N-methyl-2-(4-methyl-2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 7):-

In the same manner as in Example 5, the objective compound was prepared from methyl-(Z)-2-(4-methyl-2-phenoxyphenyl)-2-methoxyiminoacetate (Product (B)).

- 15 Example 8

Production of (E)-N-methyl-2-(4-methyl-2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 8).

- 20 In the same manner as in Example 5, the objective compound was prepared from methyl-(E)-2-(4-methyl-2-phenoxyphenyl)-2-methoxyiminoacetate.

Example 9

Production of (E)-N-methyl-2-(6-methoxy-2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 9).

- 25 In the same manner as in Example 5, the objective compound was prepared from 3-methoxydiphenyl ether.

Example 10

- 30 Production of (E)-N-methyl-2-(5-nitro-2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 10):-

- (E)-N-Methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (500 mg) was dissolved in acetic anhydride (1 ml), and the resultant mixture was cooled to 0°C. Fuming nitric acid (302 mg) was added thereto, and the mixture was stirred at the same temperature for 30 minutes. Excess methanolic ethylamine was added thereto, followed by stirring for 12 hours. The reaction mixture was diluted with water and extracted with diethyl ether. The solvent was dried and the residue was subjected to high performance liquid chromatography (HPLC) to give 86 mg of the objective compound.

In the same manner as above, there were obtained the following compounds:

- 40 (E)-N-methyl-2-[2-(4-nitrophenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 11; 167 mg);
(E)-N-methyl-2-[2-(2-nitrophenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 12; 45 mg);
(E)-N-methyl-2-(3-nitro-2-phenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 13; 43 mg).

Example 11

Production of (E)-N-methyl-2-[2-(4-chlorophenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 16):-

- 45 A solution of (E)-N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (500 mg) in dichloromethane was cooled to 0°C, and sulfuryl chloride (285 mg) was dropwise added thereto. The resultant mixture was stirred at the same temperature for 2 hours, diluted with water and extracted with diethyl ether. The solvent was dried and removed, and the residue was subjected to HPLC with a mixture of hexane and ethyl acetate to give 382 mg of the objective compound.

Example 12

Production of (E)-N-methyl-2-[2-(4-bromophenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 17):-

- 55 A solution of (E)-N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (500 mg) in carbon tetrachloride (3 ml) was cooled to 0°C, and bromine in carbon tetrachloride (1.27 mmol) was dropwise added thereto. The resultant mixture was stirred at the same temperature for 1 hour, washed with a saturated aqueous sodium hydrogencarbonate solution and extracted with dichloroethane. The solvent was dried and evaporated, and the residue was subjected to HPLC with

a mixture of hexane and ethyl acetate to give 200 mg of the objective compound.

Example 13

5 Production of (E)-N-methyl-2-(2-phenoxy-5-fluorophenyl)-2-methoxyiminoacetamide (Compound No. 18):-

In the same manner as in Example 5, the objective compound was prepared from 4-fluorodiphenyl ether.

Example 14

10 Production of (E)-N-methyl-2-[2-(4-t-butylphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 19):-

To a solution of (E)-N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (300 mg) in nitromethane (3 ml), aluminium chloride (423 mg) and t-butyl chloride (147 mg) were added, followed by stirring at room temperature for 1 hour.
15 The reaction mixture was diluted with water and extracted with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give 375 mg of the objective compound.

Example 15

20 Production of (E)-N-methyl-2-[2-(4-methoxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 22):-

(1) To a solution of 4-methoxyphenol (1.50 g) and 2-chloronitrobenzene (1.91 g) in DMF (10 ml), anhydrous potassium carbonate (3.34 g) was added, and the resultant mixture was stirred at 120°C for 12 hours. The reaction mixture was cooled to room temperature, diluted with water and extracted with diethyl ether. The solvent was dried to give 4-methoxy-2'-nitrodiphenyl ether (2.61 g).
25

¹H-NMR: 3.82 (3H, s), 6.91 (2H, d, J = 9.1), 7.02 (2H, d, J = 9.1), 7.12 (1H, t, J = 8.0), 7.45 (1H, ddd, J = 9.1, 8.0, 1.7), 7.92 (1H, dd, J = 8.0, 1.7).

(2) The thus obtained 4-methoxy-2'-nitrodiphenyl ether (2.61 g) was dissolved in a mixture of water (30 ml) and toluene (30 ml), and ammonium chloride (5.1 g) and iron powders (1.79 g) were added thereto. The resultant mixture was heated under reflux and vigorously stirred for 4 hours. The reaction mixture was cooled to room temperature, followed by removal of insoluble materials by filtration. The filtrate was extracted with diethyl ether and the solvent was dried and evaporated to give 2-(4-methoxyphenoxy)aniline (2.31 g).
30

¹H-NMR: 3.78 (3H, s), 3.80 (2H, brs), 6.65 - 6.95 (8H, m).
(3) 2-(4-Methoxyphenoxy)aniline (2.31 g) as obtained (2) above was dissolved in acetic acid (3 ml), followed by addition of conc. hydrobromic acid (3 ml). The resultant mixture was cooled to 0°C and an aqueous solution (1 ml) of sodium nitrite (741 mg) was further added thereto to make a diazonium salt, which was, without isolation, dropwise added to a solution of copper(I) bromide (918 mg) in conc. hydrobromic acid (2 ml) in another reactor. The resulting mixture was stirred at 100°C for 2 hours, cooled to room temperature and diluted with water, followed by extraction with diethyl ether. The extract was washed with a saturated aqueous sodium hydrogencarbonate solution. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give 4-methoxy-2'-bromodiphenyl ether (1.89 g).
40

¹H-NMR: 3.80 (3H, s), 6.81 - 6.98 (6H, m), 7.21 (1H, t, J = 7.3), 7.60 (1H, dd, J = 7.9, 1.6).

(4) A solution of 4-methoxy-2'-bromodiphenyl ether (1.19 g) as above obtained in tetrahydrofuran (THF) (20 ml) was cooled to -78°C, and a hexane solution (5.12 mmol) of butyl lithium and hexane was added thereto. The resultant mixture was stirred at the same temperature for 1 hour, followed by addition of dimethyl oxalate (1.01 g) in THF. The mixture was further stirred at room temperature for 2 hours. The reaction mixture was diluted with a saturated ammonium chloride solution and extracted with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give methyl 2-[2-(4-methoxyphenoxy)phenyl]-2-oxoacetate (723 mg).
50

¹H-NMR: 3.75 (3H, s), 3.82 (3H, s), 6.80 (1H, d, J = 8.3), 6.91 (2H, d, J = 9.0), 7.02 (2H, d, J = 9.0), 7.16 (1H, t, J = 7.8), 7.50 (1H, ddd, J = 8.3, 7.8, 1.7), 7.94 (1H, dd, J = 7.8, 1.7).

(5) The thus obtained methyl 2-[2-(4-methoxyphenoxy)phenyl]-2-oxoacetate was treated in the same manner as in Example 5 to give the objective compound.
55

Example 16

Production of (E)-N-methyl-2-[2-(4-trimethylsilylphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 28):-

5 In the same manner as in Example 5, the objective compound was prepared from 4-trimethylsilyl diphenyl ether.

Example 17

Production of (E)-N-methyl-2-[2-(4-iodophenoxyphenyl)-2-methoxyiminoacetamide (Compound No. 29):-

10

To a solution of (E)-N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide (300 mg) in carbon tetrachloride (10 ml), a solution of iodine monochloride (205 mg) in carbon tetrachloride (1 ml) was added, and the resultant mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with water and extracted with diethyl ether. The solvent was dried and evaporated, and the residue was subjected to HPLC with a mixture of hexane and ethyl acetate to give 269 mg of the objective compound.

15

Example 18

Production of (E)-N-methyl-2-[2-(4-chloro-3-methylphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 31):-

20

In the same manner as in Example 11, the objective compound was prepared from (E)-N-methyl-2-[2-(3-tolylloxy)phenyl]-2-methoxyiminoacetamide.

Example 19

25

Production of (E)-N-methyl-2-[2-(2-tolylloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 32):-

In the same manner as in Example 5, the objective compound was prepared from 2-phenoxytoluene.

Example 20

Production of (E)-N-methyl-2-[2-(4-tolylloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 34):-

35

In the same manner as in Example 5, the objective compound was prepared from 4-phenoxytoluene.

Example 21

Production of (E)-N-methyl-2-[2-(4-isopropylphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 36):-

40

In the same manner as in Example 14, the objective compound was prepared from 2-chloropropane and (E)-N-methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide.

Example 22

45

Production of (E)-N-methyl-2-[2-(3,4-dichlorophenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 38):-

In the same manner as in Example 15 (1), the objective compound was prepared from 3,4-dichloro-2'-bromodiphenyl ether.

Example 23

Production of (E)-N-methyl-2-[2-(3-benzyloxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 39):-

55

In the same manner as in Example 15 (1), the objective compound was prepared from 3-benzyloxy-2'-bromodiphenyl ether.

Example 24

Production of (E)-N-methyl-2-[2-(3-phenoxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 41):-

- 5 In the same manner as in Example 15 (1), the objective compound was prepared from 2-bromo-(3-phenoxyphenoxy)benzene.

Example 25

- 10 Production of (E)-N-methyl-2-[2-(4-phenoxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 42):-

In the same manner as in Example 15 (1), the objective compound was prepared from 2-bromo-(4-phenoxyphenoxy)benzene.

15 Example 26

Production of (E)-N-methyl-2-(benzyloxyphenyl)-2-methoxyiminoacetamide (Compound No. 45):-

- 20 (1) To a solution of tetrahydropyran-2-yl-phenyl ether (4.19 g) in THF (60 ml), a hexane solution (28.2 mmol) of butyl lithium was added at 0°C, and the resultant mixture was stirred at the same temperature for 2 hours and added to a solution of dimethyl oxalate (5.55 g) in THF (40 ml), followed by stirring for 12 hours. The reaction mixture was neutralized with 1N hydrochloric acid and extracted with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography to give methyl-2-(2-tetrahydropyran-2-yl-oxyphenyl)-2-oxoacetate ((3.41 g).

25 ¹H-NMR: 1.65 - 1.86 (6H, m), 3.63 - 3.75 (1H, m), 3.97 - 3.92 (1H, m), 3.92 (3H, s), 5.52 (1H, brs), 7.09 (1H, t, J = 8.0), 7.27 (1H, d, J = 8.0), 7.55 (1H, ddd, J = 8.0, 8.0, 1.8), 7.84 (1H, dd, J = 8.0, 1.8).

- (2) Methyl-2-(2-tetrahydropyran-2-yl-oxyphenyl)-2-oxoacetate (2.60 g) thus obtained was dissolved in methanol (5 ml), and O-methylhydroxylamine hydrochloride (1.70 g) was added thereto. The resultant mixture was heated under reflux for 12 hours. The solvent was removed and the residue was diluted with water and extracted with diethyl ether. Removal of the solvent on drying gave a geometric isomeric mixture of methyl 2-(2-hydroxyphenyl)-2-methoxyiminoacetate, which was then dissolved in DMF (5 ml), followed by addition of anhydrous potassium carbonate (5.26 g) and benzyl bromide (4.34 g) in order. The resultant mixture was stirred at room temperature for 12 hours, diluted with water and extracted with diethyl ether. The solvent was dried and removed, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give (Z)-N-methyl-2-(2-benzyloxyphenyl)-2-methoxyiminoacetate (835 mg). The structure of this compound was confirmed by an X-ray crystal analysis.

35 ¹H-NMR: 3.36 (3H, s), 4.00 (3H, s), 5.03 (2H, s), 6.94 - 7.01 (2H, m), 7.33 - 7.42 (6H, m), 7.73 (1H, dd, J = 7.8, 1.9).

- (3) (E)-N-Methyl-2-(2-benzyloxyphenyl)-2-methoxyiminoacetate thus obtained was treated in the same manner as in Example 5 to give the objective compound.

Example 27

Production of (E)-N-methyl-2-(2-phenoxyethylphenyl)-2-methoxyiminoacetamide (Compound No. 46):-

- 45 (1) Methyl-(E)-2-(2-tolyl)-2-methoxyiminoacetate (4.74 g) was dissolved in carbon tetrachloride (100 ml), and N-bromosuccinimide (4.89 g) and benzoylperoxide (554 mg) were added thereto. The resultant mixture was heated under reflux for 1 hour and cooled to room temperature. Insoluble materials were removed by filtration. On concentration of the solvent, the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give methyl-(E)-2-(2-bromomethylphenyl)-2-methoxyiminoacetate (6.98 g).

50 ¹H-NMR: 3.89 (3H, s), 4.07 (3H, s), 4.34 (2H, s), 7.13 - 7.17 (1H, m), 7.34 - 7.47 (3H, m).

- (2) To a solution of phenol (108 mg) in THF (2 ml), 60 % sodium hydride (46 mg) was added, followed by addition of methyl-(E)-2-(2-bromomethylphenyl)-2-methoxyiminoacetate as obtained in (1) above. The resultant mixture was heated under reflux for 8 hours and cooled to room temperature. With addition of excess methylaminemethanol solution, the mixture was stirred at room temperature for 12 hours and diluted with water, followed by extraction with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate to give 137 mg of the objective compound.

Example 28

Production of (E)-N-methyl-2-[2-(3-phenoxyphenoxy)methyl]phenyl]-2-methoxyiminoacetamide (Compound No. 47):-

5 In the same manner as in Example 27, the objective compound was prepared from 3-phenoxyphenol and methyl-(E)-2-(2-bromomethylphenyl)-2-methoxyiminoacetate.

The position of the substituent, steric configuration, melting point (m.p.) and NMR data of the objective compounds as obtained in the preceding examples are shown in Tables 1, 2 and 3 below.

10

15

20

25

30

35


40

45

50

55


Table 1



Compound No.	R ¹	R ²	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
1	H	H			3.87 (3H, s, CH ₃), 5.60 (1H, brs, NH), 6.33 (1H, brs, NH), 6.67-7.50 (9H, m, arom.)
2	CH ₃	CH ₃			2.77 (3H, s, N-CH ₃), 2.90 (3H, s, N-CH ₃), 3.97 (3H, s, O-CH ₃), 6.63-7.87 (9H, m, arom.)
3*)	H	CH ₃	E	65.5-66.5	2.78 (3H, d, J=5, N-CH ₃), 3.85 (3H, s, O-CH ₃), 6.53 (1H, brs, NH), 5=6.70-7.47 (9H, m, arom.)
4	H	CH ₃	Z	104.5-105.5	2.73 (3H, d, J=5, N-CH ₃), 3.93 (3H, s, O-CH ₃), 6.27 (1H, brs, NH), 6.67-7.87 (9H, m, arom.)

Note: *) Depending on the condition for crystallization, this compound includes transparent needle-like crystals (m.p., 104.5-105.5°C/hexane) and transparent prisms (m.p., 82-5-83°C/ethyl acetate-hexane).

Table 2



Compound No.	X_1, X_2, X_3	R^4, R^5	Steric configuration	m.p. ($^{\circ}\text{C}$)	$^1\text{H-NMR } \delta \text{ (CDCl}_3\text{) (ppm)}$
5	H, H, 3-Me	H, H	Z	-	2.31 (3H, s), 2.78 (3H, d, J = 4.9), 4.00 (3H, s), 6.31 (1H, brs), 6.76-6.92 (3H, m), 7.12-7.21 (2H, m), 7.29-7.37 (2H, m), 7.58 (1H, dd, J = 7.8, 2.0)
6	H, H, 3-Me	H, H	E	-	2.31 (3H, s), 2.87 (3H, d, J = 4.9), 3.91 (3H, s), 6.62 (1H, brs), 6.80-6.91 (4H, m), 7.10-7.21 (2H, m), 7.29-7.35 (2H, m)
7	H, H, H, 4-Me	H, 4-Me	Z	-	2.31 (3H, s), 2.87 (3H, d, J = 4.9), 3.97 (3H, s), 6.23 (1H, brs), 6.67 (1H, s), 6.90-7.33 (6H, m), 7.49 (1H, d, J = 7.8)

(Continued)

8	H, H, H	H, 4-Me	E	98-100	2.31 (3H, s), 2.86 (3H, d, J = 4.9), 3.90 (3H, s), 6.61 (1H, brs), 6.70 (1H, s), 6.83-7.33 (7H, m)
9	H, H, H	H, 6-OMe	E	106-108	2.80 (3H, d, J = 4.9), 3.75 (3H, s), 3.84 (3H, s), 6.45 (1H, d, J = 7.8), 6.64 (1H, d, J = 7.8), 6.77 (1H, brs), 7.00-7.06 (3H, m), 7.18-7.28 (3H, m)
10	H, H, H	H, 5-NO ₂	E	-	2.92 (3H, d, J = 4.9), 4.00 (3H, s), 6.80 (1H, brs), 6.83 (1H, d, J = 9.3), 7.10 (2H, dd, J = 7.3, 1.2), 7.23 (1H, t, J = 7.3), 7.40 (2H, t, J = 7.3), 8.16 (1H, dd, J = 9.3, 2.7), 8.22 (1H, d, J = 2.7)
11	H, H, 4-NO ₂	H, H	E	110-112	2.85 (3H, d, J = 4.9), 3.84 (3H, s), 6.66 (1H, brs), 7.05 (2H, d, J = 9.3), 7.30-7.47 (4H, m), 8.18 (2H, d, J = 9.3)
12	H, H, 2-NO ₂	H, H	E	-	2.81 (3H, d, J = 5.1), 3.90 (3H, s), 6.65 (1H, brs), 7.02 (1H, d, J = 8.81), 7.06 (1H, dd, J = 8.3, 1.4), 7.14 (1H, td, J = 8.3, 1.2), 7.26 (1H, td, J = 7.5, 1.2), 7.22-7.43 (2H, m), 7.46 (1H, td, J = 8.6, 1.7), 7.87 (1H, dd, J = 8.1, 1.7)

(Continued)

13	H, H, H	H, 3-NO ₂	E	-	2.76 (3H, d, J = 5.1), 3.84 (3H, s), 6.81 (2H, d, J = 7.8), 7.01 (1H, t, J = 7.3), 7.23 (2H, m), 7.37 (1H, t, J = 7.8), 7.53 (1H, dd, J = 7.8, 1.7)
14	H, H, 4-NO ₂	H, 5-NO ₂	E	172-174	2.82 (3H, d, J = 4.9), 4.11 (3H, s), 7.00 (1H, d, J = 9.0), 7.01 (1H, brs), 7.18 (2H, d, J = 9.0), 8.27 (2H, d, J = 9.0), 8.27 (1H, dd, J = 9.0, 2.9), 8.48 (1H, d, J = 2.9)
15	H, H, 4-Ac	H, H	E	112-113	2.56 (3H, s), 2.85 (3H, d, J = 4.9), 3.85 (3H, s), 6.63 (1H, brs), 7.01 (1H, m), 7.02 (2H, d, J = 8.8), 7.24 (1H, td, J = 7.5, 1.2), 7.37 (1H, m), 7.40 (1H, dd, J = 7.3, 2.0), 7.91 (2H, d, J = 8.8)
16	H, H, 4-Cl	H, H	E	108-109	2.87 (3H, d, J = 4.9), 3.90 (3H, s), 6.67 (1H, brs), 6.89 (1H, d, J = 8.5), 6.95 (2H, d, J = 9.0), 7.17 (1H, td, J = 8.5, 1.2), 7.26 (2H, d, J = 9.0), 7.30-7.38 (2H, m)
17	H, H, 4-Br	H, H	E	99-101	2.83 (3H, d, J = 4.9), 3.88 (3H, s), 6.70 (1H, brs), 6.89 (2H, d, J = 8.9), 7.15 (1H, t, J = 7.6), 7.31 (2H, d, J = 7.6), 7.38 (2H, d, J = 8.9)

(Continued)

18	H, H, H	H, 5-F	E	-	2.83 (3H, d, J = 5.1), 3.88 (3H, s), 6.65 (1H, brs), 6.86 (1H, m), 6.95-7.08 (5H, m), 7.24-7.32 (2H, m)
19	H, H, 4-tBu	H, H	E	-	1.31 (9H, s), 2.87 (3H, d, J = 5.1), 3.93 (3H, s), 6.65 (1H, brs), 6.88 (1H, dd, J = 8.3, 1.0), 6.94 (2H, d, J = 8.8), 7.12 (1H, td, J = 7.0, 1.0), 7.27-7.32 (2H, m), 7.31 (2H, d, J = 8.8)
20	H, H, H	H, 5-OMe	Z	-	2.75 (3H, d, J = 5.1), 3.83 (3H, s), 3.98 (3H, s), 6.15 (1H, brs), 6.85-6.94 (3H, m), 7.03 (1H, t, J = 7.3), 7.14 (1H, d, J = 2.7), 7.25-7.30 (2H, m), 7.36 (1H, s)
21	H, H, H	H, 5-OMe	E	-	2.84 (3H, d, J = 5.1), 3.80 (3H, s), 3.87 (3H, s), 6.57 (1H, brs), 6.83-6.96 (5H, m), 6.99-7.04 (1H, m), 7.23-7.29 (2H, m)
22	H, H, 4-OMe	H, H	E	118	2.88 (3H, d, J = 4.9), 3.79 (3H, s), 3.95 (3H, s), 6.70 (1H, brs), 6.79 (1H, d, J = 8.6), 6.84 (2H, d, J = 9.0), 6.98 (2H, d, J = 9.0), 7.11 (1H, td, J = 7.0, 1.0), 7.25-7.31 (2H, m)

(Continued)

23	H, H, H	H, 5-CF ₃	E	111	2.90 (3H, d, J = 5.1), 3.96 (3H, s), 6.74 (1H, brs), 6.88 (1H, d, J = 8.6), 7.06 (2H, d, J = 7.6), 7.16 (1H, t, J = 7.6), 7.36 (2H, t, J = 7.6), 7.52-7.57 (2H, m)
24	H, H, H	H, 5-CF ₃	Z	123-124	2.82 (3H, d, J = 4.9), 4.05 (3H, s), 6.68 (1H, brs), 6.85 (1H, d, J = 8.8), 7.03 (2H, d, J = 7.5), 7.17 (1H, t, J = 7.5), 7.36 (2H, t, J = 7.6), 7.54 (1H, dd, J = 8.8, 2.2)
25	H, H, H	H, 5-SiMe ₃	Z	80-82	0.27 (9H, s), 2.78 (3H, d, J = 4.9), 4.01 (3H, s), 6.39 (1H, brs), 6.83 (1H, d, J = 8.1), 7.00 (2H, d, J = 7.5), 7.09 (1H, t, J = 7.5), 7.28-7.31 (2H, m), 7.45 (1H, dd, J = 8.1, 1.7), 7.68 (1H, d, J = 1.7)
26	H, H, 4-SiMe ₃	H, H	Z	103-105	0.25 (9H, s), 2.78 (3H, d, J = 4.9), 3.99 (3H, s), 6.32 (1H, brs), 6.89 (1H, dd, J = 8.0, 1.1), 6.95 (2H, d, J = 8.5), 7.14 (1H, td, J = 7.6, 1.1), 7.33 (1H, ddd, J = 8.0, 7.6, 1.1), 7.45 (2H, d, J = 8.5), 7.60 (1H, dd, J = 7.6, 1.7)

(Continued)

27	H, H, H	H, 5-SiMe ₃	E	86-88	0.26 (9H, s), 2.87 (3H, d, J = 4.9), 3.91 (3H, s), 6.67 (1H, brs), 6.85 (1H, d, J = 8.10, 7.00-7.11 (3H, m), 7.27-7.33 (2H, m), 7.42-7.46 (2H, m)
28	H, H, 4-SiMe ₃	H, H	E	-	0.25 (9H, s), 2.84 (3H, d, J = 4.9), 3.89 (3H, s), 6.65 (1H, brs), 6.90 (1H, dd, J = 8.3, 1.5), 6.99 (2H, d, J = 8.6), 7.14 (1H, td, J = 7.9, 1.0), 7.28-7.34 (2H, m), 7.44 (2H, d, J = 8.6)
29	H, H, 4-I	H, H	E	96	2.85 (3H, d, J = 4.9), 3.88 (3H, s), 6.74 (1H, brs), 6.77 (2H, d, J = 8.8), 6.90 (1H, d, J = 8.3), 7.16 (1H, td, J = 8.0, 1.0), 7.30-7.37 (2H, m), 7.57 (2H, d, J = 8.8)
30	H, H, H	4-Me, 5-Cl	E	171-172	2.28 (3H, s), 2.85 (3H, d, J = 5.1), 3.91 (3H, s), 6.63 (1H, brs), 6.75 (1H, s), 6.99 (2H, d, J = 7.3), 7.09 (1H, t, J = 7.3), 7.25-7.33 (3H, m)
31	H, 4-Cl, 3-Me	H, H	E	-	2.32 (3H, s), 2.87 (3H, d, J = 4.4), 3.91 (3H, s), 6.65 (1H, brs), 6.79 (1H, dd, J = 8.7, 2.8), 6.85-6.89 (2H, m), 7.16 (1H, t, J = 7.4), 7.25-7.39 (3H, m)

(Continued)

32	H, H, 2-Me	H, H	E	-	2.21 (3H, s), 2.89 (3H, d, J = 5.1), 3.95 (3H, s), 6.64 (1H, brs), 6.68 (1H, d, J = 8.3), 6.91 (1H, d, J = 8.1), 7.00-7.32 (6H, m)
33	H, H, H	H, 5-Me	E	-	2.34 (3H, s), 2.87 (3H, d, J = 5.1), 3.89 (3H, s), 6.61 (1H, brs), 6.82 (1H, d, J = 8.3), 6.96-7.00 (2H, m), 7.05 (1H, t, J = 7.3), 7.11 (1H, s), 7.14-7.31 (3H, m)
34	H, H, 4-Me	H, H	E	113-115	2.29 (3H, s), 2.81 (3H, d, J = 4.9), 3.89 (3H, s), 6.71 (1H, brs), 6.82 (1H, d, J = 8.1), 6.89-6.92 (2H, m), 7.05-7.10 (3H, m), 7.23-7.31 (2H, m)
35	H, 2,4-di-(i)Pr	H, H	E	92-95	1.19 (6H, d, J = 6.8), 1.24 (6H, d, J = 7.0), 2.88 (3H, d, J = 5.1), 2.85 (1H, m), 3.20 (1H, sept, J = 6.8), 3.95 (3H, s), 6.67 (1H, brs), 6.71 (1H, d, J = 8.3), 6.82 (1H, d, J = 8.3), 6.98 (1H, dd, J = 8.3, 2.2), 7.06 (1H, td, J = 7.6, 1.0), 7.13 (1H, d, J = 2.2), 7.15-7.31 (2H, m)


(Continued)

36	H, H, 4-(i)Pr	H, H	E	-	1.23 (6H, d, J = 7.1), 2.86 (3H, d, J = 5.1), 2.87 (1H, m), 3.92 (3H, s), 6.66 (1H, brs), 6.84 (1H, dd, J = 8.3, 1.5), 6.94 (2H, d, J = 8.4), 7.26-7.32 (2H, m), 7.15 (2H, d, J = 8.4)
37	H, H, H	5,6-di-Cl	E	153	2.92 (3H, d, J = 4.9), 3.96 (3H, s), 6.70 (1H, d, J = 8.9), 6.73 (1H, brs), 7.01-7.09 (2H, m), 7.13-7.15 (1H, m), 7.29-7.33 (2H, m), 7.37 (1H, d, J = 8.9)
38	H, 3,4-di-Cl	H, H	E	80-81	2.88 (3H, d, J = 4.9), 3.89 (3H, s), 6.62 (1H, brs), 6.87 (1H, dd, J = 8.6, 2.7), 6.94 (1H, d, J = 8.3), 7.11 (1H, d, J = 2.7), 7.21-7.41 (4H, m),
39	H, H, 3-OCH ₂ Ph	H, H	E	-	2.86 (3H, d, J = 5.1), 3.90 (3H, s), 5.00 (2H, s), 6.58-6.73 (4H, m), 6.91 (1H, d, J = 8.3), 7.13-7.42 (9H, m)
40	H, H, 3-OMe	H, H	E	49-50	2.87 (3H, d, J = 5.1), 3.76 (3H, s), 3.91 (3H, s), 6.58-6.65 (4H, m), 6.93 (1H, d, J = 8.1), 7.12-7.22 (2H, m), 7.30-7.36 (2H, m)

(Continued)

41	H, H, 3-OPh	H, H	E	-	2.87 (3H, d, J = 4.9), 3.87 (3H, s), 6.64 (1H, brs), 6.70 (1H, brs), 6.72 (2H, brd, J = 7.8), 6.94 (1H, d, J = 7.8), 7.01 (2H, d, J = 7.8), 7.07-7.37 (7H, m)
42	H, H, 4-OPh	H, H	E	-	2.89 (3H, d, J = 4.9), 3.94 (3H, s), 6.69 (1H, brs), 6.88 (1H, d, J = 8.8), 6.94-7.13 (8H, m), 7.29-7.35 (4H, m)
43	5-OPh, 2,4-di-Cl	H, H	E	-	2.87 (3H, d, J = 5.1), 3.85 (3H, s), 6.65 (1H, brs), 6.70 (1H, s), 6.80 (1H, d, J = 8.3), 6.90 (2H, brd, J = 7.8), 7.07 (1H, brt, J = 7.5)

Table 3



Compound No.	X ₁	R ⁴	Z	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
44	H	H	-CH ₂ O-	Z	110-111	2.65 (3H, d, J = 4.9), 3.99 (3H, s), 5.03 (2H, s), 6.26 (1H, brs), 6.94-7.03 (2H, m), 7.34-7.40 (6H, m), 7.50 (1H, dd, J = 7.8, 2.0)
45	H	H	-CH ₂ O-	E	123-126	2.87 (3H, d, J = 4.9), 3.93 (3H, s), 5.06 (2H, s), 6.61 (1H, brs), 6.97-7.04 (2H, m), 7.28-7.39 (7H, m)
46	H	H	-OCH ₂ -	E	119-120	2.89 (3H, d, J = 4.9), 3.93 (3H, s), 4.95 (2H, s), 6.69 (1H, brs), 6.88-6.96 (3H, m), 7.20-7.29 (3H, m), 7.34-7.43 (2H, m), 7.53-7.54 (1H, m)

(Continued)

47	3-OPh	H	-OCH ₂ -	E	-	2.83 (3H, d, J = 4.9), 3.88 (3H, s), 4.91 (2H, s), 6.55-6.64 (3H, m), 6.70 (1H, brs), 6.98-7.02 (2H, m), 7.09 (1H, t, J = 7.3), 7.15-7.19 (2H, m), 7.21-7.42 (4H, m), 7.47-7.50 (1H, m)
48	4-Br	H	-OCH ₂ -	E	137-138	2.89 (3H, d, J = 5.1), 3.92 (3H, s), 4.92 (2H, s), 6.76 (2H, d, J = 8.8), 7.18-7.21 (1H, m), 7.34 (2H, d, J = 8.8), 7.36-8.00 (3H, m)
49	3-OMe	H	-OCH ₂ -	E	-	2.87 (3H, d, J = 4.9), 3.75 (3H, s), 3.92 (3H, s), 4.92 (2H, s), 6.45-6.51 (3H, m), 6.70 (1H, brs), 7.13 (1H, d, J = 8.1), 7.18-7.24 (1H, m), 7.33-7.43 (2H, m), 7.49-7.52 (1H, m)

Example 29

Production of (E)-N-methyl-2-[2-(3-chlorophenoxymethyl)phenyl]-2-methoxyiminoacetamide (Compound No. 50):-

5 Methyl-(E)-2-(bromomethylphenyl)-2-methoxyiminoacetate as obtained in Example 27 (1) (250 mg), 3-chlorophenol (225 mg) and potassium carbonate (482 mg) were dissolved in DMF, and the resultant mixture was stirred at room temperature for 10 hours. A solution (677 mg) of methylamine in 30 % methanol was added thereto, and stirring was continued for 10 hours. The reaction mixture was neutralized with dilute hydrochloric acid, extracted with diethyl ether. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography to give 265
10 mg of the objective compound.

In the same manner as above, Compounds Nos. 50 to 70 as shown in Table 4 below were obtained.

15

20

25

30

35


40

45

50

55

Table 4



Compound No.	Z	X ₁ , X ₂	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
50	-OCH ₂ -	H, 3-Cl	E	oily	2.89 (3H, d, J = 4.9), 3.93 (3H, s), 4.93 (2H, s), 6.74 (1H, brs), 6.78 (1H, m), 6.89-6.93 (2H, m), 7.15 (1H, d, J = 7.8), 7.22 (1H, m), 7.35-7.50 (3H, m)
51	-OCH ₂ -	H, 2-Cl	E	126	2.87 (3H, d, J = 5.1), 3.93 (3H, s), 5.05 (2H, s), 6.74 (1H, brs), 6.84-6.89 (2H, m), 7.14 (1H, dt, J = 1.5, 7.9), 7.22 (1H, dd, J = 7.3, 1.7), 7.33-7.44 (3H, m), 7.56 (1H, brd, J = 7.6)
52	-OCH ₂ -	H, 4-Ph-CH=CH-	E	215-216	2.90 (3H, d, J = 4.9), 3.94 (3H, s), 4.97 (2H, s), 6.71 (1H, brs), 6.88 (2H, d, J = 8.8), 6.95 (1H, d, J = 16.3), 7.05 (1H, d, J = 16.3), 7.20-7.54 (11H, m)

(Continued)

53	-OCH ₂ -	H, 4-Ph-CO-	E	oily	2.90 (3H, d, J = 5.1), 3.94 (3H, s), 5.04 (2H, s), 6.80 (1H, brs), 6.95 (2H, d, J = 8.8), 7.22 (1H, m), 7.41-7.56 (6H, m), 7.72-7.75 (2H, m), 7.80 (2H, d, J = 8.8)
54	-OCH ₂ -	H, 4-tBu	E	114	1.29 (9H, s), 2.88 (3H, d, J = 5.1), 3.93 (3H, s), 4.92 (2H, s), 6.65 (1H, brs), 6.82 (2H, d, J = 8.7), 7.21 (1H, m), 7.27 (2H, d, J = 8.7), 7.36-7.41 (2H, m), 7.52 (1H, m)
55	-OCH ₂	H, 4-Me	E	128-129	2.27 (3H, s), 2.88 (3H, d, J = 4.9), 3.92 (3H, s), 4.91 (2H, s), 6.69 (1H, brs), 6.78 (2H, d, J = 8.4), 7.05 (2H, d, J = 8.4), 7.21 (1H, m), 7.35-7.42 (2H, m), 7.50 (1H, m)
56	-OCH ₂	H, 4-CN	E	162	2.89 (3H, d, J = 5.1), 3.92 (3H, s), 5.01 (2H, s), 6.83 (1H, brs), 6.94 (2H, d, J = 8.3), 7.20 (1H, m), 7.37-7.48 (3H, m), 7.54 (2H, d, J = 8.3)
57	-OCH ₂ -	H, 2,4-di-Cl	E	123	2.87 (3H, d, J = 5.1), 3.92 (3H, s), 5.03 (2H, s), 6.77 (1H, brs), 6.78 (1H, d, J = 8.9), 7.09 (1H, dd, J = 8.9, 2.7), 7.20 (1H, dd, J = 7.0, 1.7), 7.33-7.40 (3H, m), 7.51 (1H, d, J = 6.8)

(Continued)

58	-OCH ₂ -	3,4-di-Cl	E	124- 126	2.89 (3H, d, J = 5.1), 3.93 (3H, s), 4.92 (2H, s), 6.74 (1H, dd, J = 9.0, 2.8), 6.76 (1H, brs), 6.98 (1H, d, J = 2.8), 7.18 (1H, m), 7.28 (1H, d, J = 9.0), 7.36-7.46 (3H, m)
59	-OCH ₂ -	3,5-di-Cl	E	102- 104	2.92 (3H, d, J = 5.1), 3.95 (3H, s), 4.92 (2H, s), 6.75 (1H, brs), 6.79 (2H, d, J = 1.7), 6.94 (1H, t, J = 1.7), 7.20 (1H, m), 7.38-7.47 (3H, m)
60	-OCH ₂ -	H, 3-CF ₃	E	69- 71	2.89 (3H, d, J = 5.1), 3.92 (3H, s), 4.99 (2H, s), 6.72 (1H, brs), 7.05 (1H, dd, J = 8.3, 2.0), 7.13 (1H, brs), 7.17-7.25 (2H, m), 7.33-7.51 (4H, m)
61	-OCH ₂ -	H, 3-Me ₂ N	E	oily	2.88 (3H, d, J = 4.9), 2.91 (6H, s), 3.93 (3H, s), 4.92 (2H, s), 6.26-6.28 (2H, m), 6.35 (1H, dd, J = 8.4, 2.2), 6.60 (1H, brs), 7.11 (1H, t, J = 8.4), 7.20 (1H, dd, J = 9.0, 2.0), 7.37-7.41 (2H, m), 7.52 (1H, dd, J = 7.0, 2.0)

(Continued)

62	-OCH ₂ -	H, 3-Br	E	56-58	2.90 (3H, d, J = 5.1), 3.94 (3H, s), 4.93 (2H, s), 6.72 (1H, brs), 6.82 (1H, dt, J = 7.6, 2.0), 7.05-7.14 (3H, m), 7.20 (1H, m), 7.35-7.50 (3H, m)
63	-OCH ₂ -	H, 3-NO ₂	E	127-129	2.92 (3H, d, J = 4.9), 3.95 (3H, s), 5.05 (2H, s), 6.79 (1H, brs), 7.20-7.23 (2H, m), 7.37-7.43 (3H, m), 7.49 (1H, m), 7.73 (1H, t, J = 2.3), 7.81 (1H, dd, J = 8.0, 2.3)
64	-OCH ₂ -	H, 4-F	E	97	2.89 (3H, d, J = 5.1), 3.93 (3H, s), 4.91 (2H, s), 6.73 (1H, brs), 6.79-6.84 (2H, m), 6.90-6.97 (2H, m), 7.20 (1H, m), 7.34-7.43 (2H, m), 7.50 (1H, m)
65	-OCH ₂ -	H, 3-(i)PrO	E	oily	1.31 (6H, d, J = 6.1), 2.88 (3H, d, J = 4.9), 3.93 (3H, s), 4.50 (1H, sept, J = 6.1), 4.91 (2H, s), 6.44-6.50 (3H, m), 6.68 (1H, brs), 7.10-7.22 (2H, m), 7.36-7.4 (2H, m), 7.51 (1H, d, J = 6.8)

(Continued)

66	-OCH ₂ -	H, 4-Cl	E	132-133	2.90 (3H, d, J = 4.9), 3.93 (3H, s), 4.92 (2H, s), 6.72 (1H, brs), 6.81 (2H, d, J = 9.0), 7.20 (1H, m), 7.20 (2H, d, J = 9.0), 7.37-7.49 (3H, m)
67	-SCH ₂ -	H, H	E	oily	2.90 (3H, d, J = 5.1), 3.95 (3H, s), 3.99 (2H, s), 6.75 (1H, brs), 7.14-7.31 (9H, m)
68	-SOCH ₂ -	H, H	E	oily	2.94 (3H, d, J = 4.9), 3.76 (1H, d, J = 12.5), 3.98 (3H, s), 4.24 (1H, d, J = 12.5), 6.73 (1H, d, J = 7.8), 7.03 (1H, brs), 7.14-7.20 (2H, m), 7.30-7.45 (6H, m)
69	-OCH ₂ -	H, 4-Cl	Z	162-163	2.90 (3H, d, J = 4.9), 3.98 (3H, s), 5.17 (2H, s), 6.64 (1H, brs), 6.85 (2H, brd, J = 8.9), 7.23 (2H, brd, J = 8.9), 7.36-7.52 (4H, m)
70	-OCH ₂ -	H, 3-(i)Pro	Z	oily	1.32 (6H, d, J = 6.1), 2.87 (3H, d, J = 4.9), 3.95 (3H, s), 4.50 (1H, sept, J = 6.1), 5.16 (2H, s), 6.49-6.52 (2H, m), 6.67 (1H, brs), 7.12-7.54 (6H, m)

Example 30

Production of N-methyl-2-[2-(3-methyl-2-butenyloxy)phenyl]-2-methoxyiminoacetamide (Compound No. 71):-

(1) A solution of 2-bromophenol (10.0 g), dihydropyran (7.28 g) and pyridinium p-toluenesulfonate (PPTS) (725 mg) in dichloromethane was stirred at room temperature for 10 hours. After removal of the solvent and excess dihydropyran, the reaction mixture was diluted with THF (150 ml), followed by cooling to -78°C. n-Butyl lithium (1.6M; 47 ml) was dropwise added thereto, and the mixture was stirring for 30 minutes. Dimethyl oxalate (13.6 g) was added to the mixture at the same temperature and stirring was continued for 2 hours. The reaction mixture was diluted with water and extracted with ether, followed by removal of the solvent on drying. The residue was dissolved in methanol (100 ml) and combined with O-methyl hydroxylamine hydrochloride (7.24 g), followed by heating under reflux for 4 hours. The reaction mixture was allowed to cool to room temperature and, after removal of methanol, diluted with water, followed by extraction with ethyl acetate. The solvent was dried and distilled off, and the residue was dissolved in 30 % methanolic methylamine (23 g). The resultant mixture was stirred for 10 hours and methylamine was removed therefrom. The residue was purified by silica gel column chromatography, whereby 3.23 g of E-isomer and 6.87 g of Z-isomer of N-methyl-[2-(2-hydroxy)phenyl]-2-methoxyiminoacetamide were obtained.


E-isomer:-

¹H-NMR: 3.90 (3H, s), 4.15 (3H, s), 6.14 (1H, brs), 6.96 - 7.01 (2H, m), 7.24 (1H, dd, J = 8.3, 1.7), 7.35 (1H, brt, J = 8.3).

(2) The above obtained E-isomer (200 mg), prenyl bromide (286 mg) and potassium carbonate (265 mg) was dissolved in DMF, and the resultant solution was stirred at room temperature for 2 hours. The reaction mixture was neutralized with dilute hydrochloric acid, extracted with ethyl acetate and purified by silica gel column chromatography to give 286 mg of the objective compound.

In the same manner as above, Compounds Nos. 71 to 74 as shown in Table 5 below were obtained.

Table 5



Compound No.	A	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
71	$\begin{array}{c} \text{CH}_3 \diagup \\ \text{C}=\text{CHCH}_2- \\ \text{CH}_3 \diagdown \end{array}$	E	48-50	1.69 (3H, s), 1.75 (3H, s), 2.90 (3H, d, J = 4.9), 3.94 (3H, s), 4.50 (2H, d, J = 6.6), 5.37 (1H, t, J = 6.6), 6.64 (1H, brs), 6.94 (1H, dd, J = 8.1, 1.0), 6.99 (1H, dd, J = 7.6, 1.0), 7.23 (1H, dd, J = 7.6, 1.7), 7.33 (1H, ddd, J = 8.1, 7.6, 1.7)
72	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 \diagup \text{C}=\text{CH}(\text{CH}_2)_2 \text{C}=\text{CHCH}_2- \\ \text{CH}_3 \diagdown \end{array}$	E	oily	1.60 (3H, s), 1.68 (6H, s), 2.91 (3H, d, J = 4.9), 3.94 (3H, s), 4.53 (2H, d, J = 5.1), 5.08 (1H, t, J = 5.1), 5.38 (1H, t, J = 5.1), 6.64 (1H, brs), 6.94 (1H, d, J = 8.3), 6.99 (1H, dd, J = 7.6, 1.0), 7.23 (1H, dd, J = 7.6, 1.7), 7.34 (1H, ddd, J = 8.3, 7.6, 1.7)

(Continued)

73	$\begin{array}{c} \text{Cl} \diagup \\ \text{C}=\text{CHCH}_2- \\ \text{Cl} \diagdown \end{array}$	E	41	2.93 (3H, d, J = 4.9), 3.96 (3H, s), 4.66 (2H, d, J = 6.1), 6.07 (1H, t, J = 6.1), 6.71 (1H, brs), 6.91 (1H, brd, J = 8.3), 7.04 (1H, brt, J = 8.3), 7.23 (1H, dd, J = 7.6, 1.7), 7.38 (1H, ddd, J = 8.3, 7.6, 1.7),
74	$\begin{array}{c} \text{Cl} \diagup \\ \text{C}=\text{CHCH}_2- \\ \text{Cl} \diagdown \end{array}$	Z	112	2.96 (3H, d, J = 4.9), 4.02 (3H, s), 6.87 (1H, d, J = 7.9), 7.02 (1H, t, J = 7.6), 7.37 (1H, td, J = 7.9, 1.7), 7.49 (1H, dd, J = 7.6, 1.7)

Example 31

Production of (E)-N-methyl-2-[2-(pyrimidin-2-yloxymethyl)phenyl]-2-methoxyiminoacetamide (Compound No. 79):-

(1) Methyl-2-(2-bromomethylphenyl)-2-methoxyiminoacetate (2.00 g) and potassium acetate (1.37 g) were suspended in THF, and the suspension was heated under reflux for 3 hours. The reaction mixture was cooled to room temperature, and 30 % methanolic methylamine (5.4 g) was added thereto. The resultant mixture was stirred for 10 hours, diluted with water and neutralized, followed by extraction with ethyl acetate. The solvent was dried and distilled off, and the residue was purified by silica gel column chromatography, whereby there was obtained (E)-N-methyl-2-(2-hydroxymethylphenyl)-2-methoxyiminoacetamide (1.05 g).

¹H-NMR: 2.92 (3H, d, J = 4.9), 3.22 (1H, t, J = 4.9), 3.95 (3H, s), 4.39 (2H, d, J = 4.9), 6.97 (1H, brs), 7.13 (1H, brd, J = 7.6), 7.35 (1H, td, J = 7.6, 1.5), 7.42 (1H, td, J = 7.6, 1.5), 7.52 (1H, brd, J = 7.6).

(2) (E)-N-methyl-2-(2-hydroxymethylphenyl)-2-methoxyiminoacetamide as obtained in (1) above (300 mg), 2-chloropyrimidine (233 mg) and sodium hydride (40 %; 135 mg) were dissolved in DMF, and the resultant mixture was stirred at 100°C for 6 hours. After neutralization, the reaction mixture was diluted with water and extracted with dichloromethane. The solvent was dried and evaporated, and the residue was purified by silica gel column chromatography to give 70 mg of the objective compound.

Example 32

Production of N-methyl-2-(2-phenylthiophenyl)-2-methoxyiminoacetamide (Compound No. 82):-

The objective compound was prepared in the same manner as in Example 5 but using phenylsulfide in place of 3-phenoxytoluene.

Example 33

Production of (E)-N-methyl-2-(2-phenylsulfinylmethylphenyl)-2-methoxyiminoacetamide (Compound No. 68):-

The objective compound was produced by oxidation of Compound No. 67 with sodium metaperiodate.

Example 34

Production of (E)-N-methyl-2-(5-trifluoromethylpyridyl-2-oxyphenyl)-2-methoxyiminoacetamide (Compound No. 75):-

(1) To a solution of 2-bromophenol (3.46 g) in DMF (10 ml), potassium carbonate (5.53 g) and 2-chloro-5-trifluoromethyl pyridine (7.26 g) were added, and the resultant mixture was stirred at 60°C, followed by cooling. Water was added to the reaction mixture, which was extracted with dichloromethane. The extract was dried over anhydrous sodium sulfate and purified by silica gel column chromatography to give 2-(5-trifluoromethylpyridyl-2-oxy-bromobenzene (2.70 g).


(2) 2-(5-trifluoromethylpyridyl-2-oxy)-bromobenzene obtained in (1) above (2.62 g) was dissolved in THF (20 ml). Under argon stream, 1.6 M hexane solution (6.2 ml) of n-butyl lithium was dropwise added thereto at -78°C in 2 minutes, and the resultant mixture was stirred for 5 minutes. A solution of dimethyl oxalate (1.95 g) in THF (20 ml) was added thereto at once, and thereafter warmed to room temperature, followed by stirring for 20 minutes. Aqueous ammonium chloride was added to the reaction mixture and, after removal of THF, extraction with dichloromethane was performed. The residue was washed with water and dried over anhydrous sodium sulfate. After removal of the solvent, the residue was purified by silica gel column chromatography to afford methyl-2-(5-trifluoromethylpyridyl-2-oxyphenyl)-2-oxoacetate (1.68 g).

(3) Methyl-2-(5-trifluoromethylpyridyl-2-oxyphenyl)-2-oxoacetate (1.21 g) as obtained (2) was dissolved in methanol (10 ml), and O-methyl hydroxylaminehydrochloride (0.74 g) was added thereto. The resultant mixture was refluxed for 2 hours, followed by removal of methanol. Water was added to the mixture, which was extracted with dichloromethane. The residue was dried over anhydrous sodium sulfate and, after removal of the solvent, purified by silica gel column chromatography to give methyl-2-(5-trifluoromethylpyridyl-2-oxyphenyl)-2-methoxyiminoacetate in a mixture of Z-isomer (0.2 g) and E-isomer (0.60 g).

(4) The E-isomer (0.60 g) was dissolved in methanol (10 ml), and 30 % methanolic methyl amine (0.35 g) was dropwise added thereto. The resultant mixture was refluxed for 20 minutes, and after removal of methanol, the residue was purified by silica gel column chromatography and recrystallized from n-hexane to give 0.50 g of the objective compound.

In the same manner as above, Compounds No. 75 to 77 as shown in Table 6 below were obtained.

Table 6



Compound No.	X_1	Steric configuration	m.p. ($^{\circ}C$)	1H -NMR δ ($CDCl_3$) (ppm)
75	5- CF_3	E	100-100.5	2.83 (3H, d, $J = 4.9$), 3.78 (3H, s), 6.59 (1H, brs), 6.97 (1H, d, $J = 8.8$), 7.20-7.50 (4H, m), 7.87 (1H, dd, $J = 8.8, 2.9$), 8.43 (1H, d, $J = 2.9$)
76	3- CF_3	E	134.5-135.5	2.84 (3H, d, $J = 4.9$), 3.74 (3H, s), 6.58 (1H, brs), 7.06 (1H, m), 7.26-7.51 (4H, m), 7.96 (1H, d, $J = 6.4$), 8.25 (1H, d, $J = 3.7$)
77	5- CF_3	Z	85.5-86.5	2.75 (3H, d, $J = 4.9$), 3.91 (3H, s), 6.25 (1H, brs), 6.99 (1H, d, $J = 8.8$), 7.15 (1H, d, $J = 7.8$), 7.30 (1H, m), 7.47 (1H, m), 7.66 (1H, dd, $J = 7.8, 2.0$), 7.90 (1H, dd, $J = 8.8, 2.7$), 8.42 (1H, brs)


Example 35




Production of (E)-N-methyl-2-(5-trifluoromethylpyridyl-2-oxymethylphenyl)-2-methoxyiminoacetamide (Compound No. 78):-

To a solution of (E)-N-methyl-2-(2-hydroxymethylphenyl)-2-methoxyiminoacetamide (0.10 g) in THF (10 ml), sodium hydride (60 % oily suspension) (0.02 g) was added thereto, and the resultant mixture was stirred at room temperature for 10 minutes. To the reaction mixture, 2-chloro-5-trifluoromethylpyridine (0.10 g) was added, and stirring was continued at 60°C for 1 hour. The reaction mixture was cooled, diluted with water, followed by extraction with dichloromethane. The solvent was dried over anhydrous sodium sulfate, and after removal of the solvent, purified by silica gel column chromatography and recrystallized from n-hexane to give 0.11 g of the objective compound.

In the same manner as above, Compounds Nos. 78 to 81 as shown in Table 7 were obtained:

Table 7



Compound No.	A	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
78		E	85.6-86	2.89 (3H, d, J = 4.9), 3.92 (3H, s), 5.30 (2H, s), 6.76 (1H, brs), 6.78 (1H, d, J = 8.8), 7.21 (1H, m), 7.35-7.41 (2H, m), 7.53 (1H, m), 7.74 (1H, dd, J = 8.8, 2.9), 8.38 (1H, d, J = 2.9)
79		E	oily	2.93 (3H, d, J = 4.9), 3.92 (3H, s), 5.31 (2H, s), 6.90 (1H, t, J = 4.6), 6.92 (1H, brs), 7.25 (1H, dd, J = 7.5, 2.0), 7.33-7.42 (2H, m), 7.61 (1H, dd, J = 8.0, 2.0), 8.45 (2H, d, J = 4.6)
80		E	74-75	2.89 (3H, d, J = 5.1), 3.92 (3H, s), 5.24 (2H, s), 6.71 (1H, d, J = 8.1), 6.72 (1H, brs), 6.85 (1H, dd, J = 6.1, 5.1), 7.23 (1H, m), 7.36-7.41 (2H, m), 7.51-7.58 (2H, m), 8.11 (1H, dd, J = 5.1, 2.0)

(Continued)

2.90 (3H, d, J = 4.9), 3.91 (3H, s),
5.50 (2H, s), 6.77 (1H, d, J = 8.3),
6.86 (1H, m), 6.95 (1H, brs), 7.32-
7.41 (2H, m), 7.47-7.62 (3H, m),
8.10 (1H, dd, J = 5.1, 2.0)

107-108

Z



81

Example 36

Production of (E)-N-methyl-2-[2-(3-hydroxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 83):-

(1) In the same manner as in Example 15, 3-(2-bromophenoxy)phenol was prepared from resorcinol and 2-chloronitrobenzene.

(2) To a solution of 3-(2-bromophenoxy)phenol as obtained above (6.12 g) in dichloromethane (120 ml), dihydropyran (3.16 ml) and PPTS (0.1 g) were added, and the resultant mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with dichloromethane and washed with saturated sodium hydrogen carbonate solution. The solvent was dried and removed, and the residue was purified by silica gel chromatography to give 3-(2-bromophenoxy)phenol tetrahydropyranyl ether (7.18 g).

(3) A solution of 3-(2-bromophenoxy)phenyl tetrahydropyranyl ether obtained in (2) above (7.18 g) in THF (120 ml) was cooled to -78°C, and a solution of butyl lithium-hexane (1.6 M, 12.8 ml) was dropwise added thereto. The resultant mixture was stirred at the same temperature for 30 minutes, and a solution of dimethyl oxalate (4.86 g) in THF (40 ml) was added thereto. Stirring was continued at room temperature for 2 hours. The reaction mixture was diluted with water and extracted with ether. The solvent was dried and removed, and the residue was dissolved in methanol (100 ml). O-Methyl hydroxylamine hydrochloride (2.58 g) was added to the solution, which was refluxed for 3 hours. The reaction mixture was cooled to room temperature and methanol was removed therefrom, followed by dilution with water. The dilute mixture was extracted with ethyl acetate. The solvent was removed and the residue was purified by silica gel column chromatography to give methyl-2-[2-(3-hydroxyphenoxy)phenyl]-2-methoxyiminoacetate in a mixture of Z-isomer (1.48 g) and E-isomer (2.57 g).

(4) The product as obtained above was treated in the same manner as in Example 5 (3) to give the objective compound.

Example 37

Production of (E)-N-methyl-2-[2-(3-tetrahydropyran-2-yloxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 84):-

To a solution of methyl-2-[2-(3-hydroxyphenoxy)phenyl]-2-methoxyiminoacetate (0.25 g) in dichloromethane (10 ml), dihydropyran (0.3 ml) and PPTS (20 mg) were added, and the resultant mixture was stirred at room temperature for 2 hours. The reaction mixture was diluted with dichloromethane, washed with a saturated aqueous sodium hydrogencarbonate solution, followed by removal of the solvent. The residue was dissolved in methanol (3 ml), and 40 % methanolic methylamine (0.5 ml) was added thereto. The resultant mixture was stirred at room temperature for 15 hours. The solvent and excess methylamine were removed, and the residue was purified by silica gel column chromatography to give 0.27 g of the objective compound.

Example 38

Production of (E)-N-methyl-2-[2-[3-(1,3-benzothiazol-2-yloxy)phenoxy]phenyl]-2-methoxyiminoacetamide (Compound No. 90):-

To a solution of (E)-N-methyl-2-[2-(3-hydroxyphenoxy)phenyl]-2-methoxyiminoacetamide (0.21 g) in DMF (3 ml), sodium hydride (34 mg) and 2-chloro-1,3-benzothiazole (0.18 g) were added, and the resultant mixture was stirred at 60°C for 1 hour. The reaction mixture was cooled to room temperature, diluted with water and extracted with ether. The solvent was dried and the residue was purified by silica gel column chromatography to give 0.27 g of the objective compound.

Example 39

Production of (E)-N-methyl-2-[2-(3-benzoyloxyphenoxy)phenyl]-2-methoxyiminoacetamide (Compound No. 94):-

To a solution of (E)-N-methyl-2-[2-(3-hydroxyphenoxy)phenyl]-2-methoxyiminoacetamide (0.21 g) in dichloromethane (5 ml), benzoyl chloride (0.12 g) and triethylamine (0.20 g) were added, and the resultant mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with ether, washed with dilute hydrochloric acid and a saturated aqueous sodium hydrogen carbonate solution. The solvent was dried and removed, and the residue was purified by silica gel column chromatography to give 0.27 g of the objective compound.


Example 40


Production of (E)-N-methyl-2-{2-[3-(3-methoxyphenoxy)phenoxy]phenyl}-2-methoxyiminoacetamide (Compound No. 87):-

In the same manner as in Example 15, the objective compound was prepared from 3-(3-methoxyphenoxy)phenol and 2-chloronitrobenzene.





In the same manner as above, Compounds Nos. 82 to 102 as shown in Table 8 below were obtained.

Table 8







Compound No.	Z	X_1	Steric configuration	m.p. ($^{\circ}C$)	1H -NMR ($CDCl_3$) (ppm)
82	S	H	E	105-107	2.90 (3H, d, $J = 5.1$), 3.87 (3H, s), 6.69 (1H, brs), 7.17-7.41 (9H, m)
83	O	3-OH	E	157-159	2.84 (3H, d, $J = 5.9$), 3.91 (3H, s), 6.42 (1H, t, $J = 2.0$), 6.48-6.53 (2H, m), 6.75 (2H, brs), 6.90 (1H, d, $J = 8.8$), 7.05-7.33 (4H, m)
84	O	3-O- 	E	oil	1.59-2.04 (6H, m), 2.87 (3H, d, $J = 4.90$), 3.58 (1H, brd), 3.90 (1H), 3.91 (3H, s), 5.36 (1H, t, $J = 3.9$), 6.61-6.80 (5H, m), 6.93 (1H, d, $J = 7.8$), 7.15-7.35 (4H, m)


(Continued)

85	O		E	oily	2.84 (3H, d, J = 4.9), 3.87 (1H, s), 6.69 (1H, brs), 6.84-7.35 (9H, m), 8.56 (2H, d, J = 4.9)
86	O		E	110-113	2.85 (3H, d, J = 4.9), 3.87 (3H, s), 6.71 (1H, brs), 6.81-7.03 (5H, m), 7.18 (1H, brt, J = 6.8), 7.30-7.40 (3H, m), 7.90 (1H, dd, J = 2.0, 8.8), 8.43 (1H, brs)
87	O		E	oily	2.88 (3H, d, J = 4.9), 3.78 (3H, s), 3.87 (3H, s), 6.57-6.75 (7H, m), 6.95 (1H, d, J = 8.8), 7.15-7.34 (5H, m)
88	O		E	oily	2.87 (3H, d, J = 4.9), 3.87 (3H, s), 6.64-6.79 (4H, m), 6.98 (1H, d, J = 7.8), 7.10 (1H, d, J = 7.8), 7.15-7.40 (5H, m), 7.53 (1H, t, J = 8.8), 7.95 (1H, d, J = 6.8)



(Continued)

89	O		E	oily	2.75 (3H, d, J = 4.9), 3.82 (3H, s), 6.77 (1H, d, J = 8.8), 6.94-7.46 (10H, m), 7.63 (1H, t, J = 6.8), 6.77 (1H, d, J = 7.8), 7.85 (1H, d, J = 8.8), 8.14 (1H, J = 8.8)
90	O		E	oily	2.84 (3H, d, J = 4.9), 3.86 (3H, s), 6.77 (1H, brs), 6.89 (1H, dd, J = 2.0, 8.8), 7.03-7.43 (9H, m), 7.69 (1H, d, J = 8.8), 7.73 (1H, d, J = 8.8)
91	O		E	oily	2.84 (3H, d, J = 4.9), 3.88 (3H, s), 5.25 (2H, s), 6.59-6.71 (4H, m), 6.94 (1H, d, J = 7.8), 7.12-7.62 (7H, m), 7.98 (2H, d, J = 8.8)
92	O		E	oily	2.45 (3H, s), 2.86 (3H, d, J = 4.9), 3.85 (3H, s), 6.66-6.89 (5H, m), 7.15-7.34 (6H, m), 7.72 (2H, d, J = 8.8)

(Continued)

93	O	3-OCOCH ₃	E	97-98.5	2.26 (3H, s), 2.85 (3H, d, J = 4.9), 3.88 (3H, s), 6.65 (1H, brs), 6.73-6.88 (3H, m), 6.98 (1H, d, J = 8.8), 7.15-7.38 (4H, m)
94	O	3-OCOPh	E	oily	2.87 (3H, d, J = 5.8), 3.89 (3H, s), 6.69 (1H, brs), 6.88-7.04 (4H, m), 7.16-7.40 (5H, m), 7.50 (2H, t, J = 7.8), 7.63 (1H, t, J = 7.9), 8.17 (2H, d, J = 6.8)
95	O	3-OCH ₂ - 	E	oily	2.87 (3H, d, J = 4.9), 3.91 (3H, s), 5.03 (2H, s), 6.61-6.71 (4H, m), 6.91 (1H, d, J = 8.8), 7.14-7.71 (8H, m)
96	O	3-OCH ₂ C≡CH	E	80-81	2.50 (1H, t, J = 2.0), 2.87 (3H, d, J = 5.9), 3.91 (3H, s), 4.64 (2H, d, J = 2.0), 6.61-6.73 (4H, m), 6.95 (1H, d, J = 7.9), 7.13-7.37 (4H, m)

(Continued)

97	O	3-CH=CH-Ph	E	oily	2.85 and 2.87 (total 3H, d, J = 4.9), 3.85 and 3.91 (total 3H, s), 6.56-7.50 (16H, m)
98	O	3-CH=CH- 	E	oily	2.87 (3H, d, J = 4.9), 3.87 and 3.90 (total 3H, s), 6.27-7.39 (14H, m)
99	O	4-OH	E	198	2.89 (3H, d, J = 5.1), 3.95 (3H, s), 6.65 (1H, brs), 6.76-6.81 (3H, m), 6.91-6.95 (2H, m), 7.09 (1H, m), 7.26-7.29 (2H, m)
100	O	4-OCH ₂ Ph	E	130-131	2.88 (3H, d, J = 5.1), 3.94 (3H, s), 5.04 (2H, s), 6.65 (1H, brs), 6.80 (1H, dd, J = 8.6, 1.2), 6.92 (2H, d, J = 9.3), 7.09 (1H, brt, J = 7.1), 7.27-7.45 (7H, m)
101	O	3-O- 	Z	oily	2.78 (3H, d, J = 4.9), 3.97 (3H, s), 6.42 (1H, brs), 6.77 (1H, t, J = 2.4), 6.86 (2H, m), 7.01 (2H, m), 7.18 (1H, dt, J = 1.2, 7.6), 7.31-7.40 (2H, m), 7.62 (1H, dd, J = 1.7, 7.6), 7.90 (1H, dd, J = 2.2, 8.5), 8.41 (1H, brs)

(Continued)

2.77 (3H, d, J = 4.9), 3.99
 (3H, s), 5.01 (2h, s), 6.25
 (1H, brs), 6.56 (1H, dd, J
 = 1.7, 7.6), 6.61 (1H, t,
 J = 2.4), 6.72 (1H, dd, J
 = 1.7, 8.3), 6.88 (1H, dd,
 J = 1.2, 8.3), 7.14 (1H,
 dt, J = 1.0, 7.6), 7.19
 (1H, t, J = 8.3), 7.29-
 7.41 (6H, m), 7.60 (1H,
 dd, J = 1.7, 7.6)

oily

Z

3-OCH₂Ph

O

102

Example 41

Production of (E)-N-methyl-2-[2-(3-methylanilinomethyl)phenoxyphenyl]-2-methoxyiminoacetamide (Compound No. 104):-

(1) To a solution of (E)-methyl-2-[2-(3-methylphenoxy)phenyl]-2-methoxyiminoacetate (4.60 g) in benzene (120 ml), N-bromosuccinimide (2.88 g) and benzoyl peroxide (0.37 g) were added, and the resultant mixture was refluxed for 1 hour, followed by cooling to room temperature. Insoluble materials were removed by filtration and, after removal of the solvent, the residue was purified by silica gel column chromatography to give (E)-methyl-2-[2-(3-bromomethylphenoxy)phenyl]-2-methoxyiminoacetate (4.60 g).

(2) The above obtained product (0.30 g) was dissolved in DMF (4 ml), and N-methylaniline (0.13 ml) and potassium carbonate (0.16 g) were added thereto, followed by stirring at room temperature for 4 hours. The reaction mixture was diluted with water and extracted with ether. The solvent was dried and the residue was combined with 40 % methanolic methylamine (1 ml), followed by stirring at room temperature for 3 hours. The solvent was removed and the residue was purified by silica gel column chromatography to give 0.31 g of the objective compound.

Example 42

Production of (E)-N-methyl-2-[2-[3-(2-pyridyloxy)phenoxy]phenyl]-2-methoxyiminoacetamide (Compound No. 105):-

A solution of (E)-methyl-2-[2-(3-bromomethylphenoxy)phenyl]-2-methoxyiminoacetate (0.50 g), 2-hydroxypyridine (0.15 g) and silver carbonate (0.22 g) in hexane (10 ml) was heated under reflux for 4 hours, followed by removal of hexane. The reaction mixture was diluted with dichloromethane, and insoluble materials were removed by filtration. After removal of the solvent, the residue was combined with 40 % methanolic methylamine (2 ml) and stirred at room temperature for 4 hours. Methylamine was removed and the residue was purified by silica gel column chromatography to give 0.38 g of the objective compound.


Example 43

Production of (E)-N-methyl-2-[2-[3-pyrimidin-2-yloxymethyl)phenoxy]phenyl]-2-methoxyiminoacetamide (Compound No. 106):-

The objective compound was prepared from (E)-methyl-2-[2-(3-bromomethylphenoxy)phenyl]-2-methoxyiminoacetate in the same manner as in Example 31.




In the same manner as above, Compounds Nos. 103 to 112 as shown in Table 9 were obtained.

Table 9



Compound No.	R^7	Z'	Steric configuration	m.p. ($^{\circ}C$)	1H -NMR δ ($CDCl_3$) (ppm)
103	Ph	S	E	oily	2.85 (3H, d, $J = 4.9$), 3.88 (3H, s), 4.06 (2H, s), 6.61 (1H, brs), 6.80 (1H, d, $J = 7.8$), 6.86 (1H, brd, $J = 7.8$), 6.96 (1H, brs), 7.03 (1H, d, $J = 7.8$), 7.11-7.34 (9H, m)
104	Ph	NCH_3	E	oily	2.84 (3H, d, $J = 5.9$), 3.02 (3H, s), 3.86 (3H, s), 4.48 (2H, s), 6.59 (1H, brs), 6.72 (2H, d, $J = 7.8$), 6.85-6.97 (4H, m), 7.12-7.37 (7H, m)
105		O	E	oily	2.86 (3H, d, $J = 4.9$), 3.89 (3H, s), 5.34 (2H, s), 6.61 (1H, brs), 6.80 (1H, d, $J = 8.8$), 6.86-6.95 (3H, m), 7.14-7.35 (6H, m), 7.58 (1H, dt, $J = 2.0, 6.8$), 8.16 (1H, dd, $J = 2.0, 4.9$)

(Continued)

106	Ph	S=O	E	oily	2.86 (3H, d, J = 5.9), 3.90 (3H, s), 3.94, 4.02 (each 1H, ABq, 11.7), 6.70-6.81 (4H, m), 6.92 (1H, dd, J = 2.0, 7.8), 7.16-7.47 (9H, m)
107		O	E	oily	2.86 (3H, d, J = 5.1), 3.88 (3H, s), 5.39 (2H, s), 6.64 (1H, brs), 6.89-6.97 (3H, m), 7.12-7.37 (6H, m), 8.52 (2H, d, J = 4.6)
108		O	E	oily	2.85 (3H, d, J = 4.9), 3.76 (3H, s), 3.89 (3H, s), 4.97 (2H, s), 6.62 (1H, brs), 6.82 (2H, d, J = 9.4), 6.89 (2H, d, J = 9.4), 6.83-7.33 (8H, m)
109		O	E	oily	2.84 (3H, d, J = 4.6), 3.88 (3H, s), 4.97 (2H, s), 6.65 (1H, brs), 6.81 (2H, d, J = 8.8), 6.87 (1H, m), 6.97 (1H, d, J = 7.8), 7.06 (1H, brs), 7.15-7.18 (2H, m), 7.27-7.34 (3H, m), 7.35 (2H, d, J = 8.8)
110	Ph	O	E	oily	2.84 (3H, d, J = 4.9), 3.88 (3H, s), 5.01 (2H, s), 6.63 (1H, brs), 6.88-7.34 (13H, m)

(Continued)



111		O	Z	oily	2.70 (3H, d, J = 4.9), 3.99 (3H, s), 5.35 (2H, s), 6.28 (1H, brs), 6.79 (1H, d, J = 8.6), 6.88 (3H, m), 7.10 (1H, brs), 7.12-7.36 (4H, m), 7.58 (2H, brt), 8.15 (1H, dd, J = 2.0, 5.1)
112	Ph	O	Z	oily	2.76 (3H, d, J = 4.9), 3.99 (3H, s), 5.03 (2H, s), 6.29 (1H, brs), 6.86-6.99 (5H, m), 7.07 (1H, brs), 7.12-7.37 (6H, m), 7.60 (1H, dd, J = 1.7, 7.6)

Example 44

Production of N-methyl-2-(2-(Z)-styrylphenyl)-2-(E)-methoxyiminoacetamide (Compound No. 113):-

(1) Following the procedure in Example 15, methyl-2-(2-styrylphenyl)-2-(E)-methoxyiminoacetate was prepared from a mixture, i.e. cis- and trans-isomer, of 2-bromostilbene.

(2) To a solution of thus prepared methyl-2-(2-styrylphenyl)-2-(E)-methoxyiminoacetate (0.70 g) in methanol (2 ml), 40 % methanolic methylamine (3 ml) was added, and the resultant mixture was stirred at room temperature for 4 hours. The solvent was removed and the residue was purified by silica gel column chromatography to give 0.26 g of the objective compound, and the mixture (0.45 g) of the objective compound and its isomer, i.e. N-methyl-2-(2-(E)-styrylphenyl)-2-(E)-methoxyiminoacetamide.

Example 45

Production of N-methyl-2-(2-(E)-styrylphenyl)-2-(E)-methoxyiminoacetamide (Compound No. 114):-

To a solution of N-methyl-2-(2-(Z and E)-styrylphenyl)-2-(E)-methoxyiminoacetamide (0.19 g) in toluene (3 ml), iodine (20 g) was added, and the resultant mixture was heated under reflux for 20 hours. The solvent was removed and the residue was purified by silica gel column chromatography to give 0.19 g of the objective compound.

Example 46

Production of (E)-N-methyl-2-[2'-((1"S*,2"R*)-1",2"-epoxy-2"-phenylethyl)phenyl]-2-methoxyiminoacetamide (Compound No. 115):-

To a solution of N-methyl-2-(2-(Z)-styrylphenyl)-2-(E)-methoxyiminoacetamide (0.26 g) in dichloromethane (5 ml), m-chloroperoxybenzoic acid (0.22 g) was added, and the resultant mixture was stirred at room temperature for 3 hours. The reaction mixture was diluted with ether and washed with aqueous sodium thiosulfate. The solvent was dried and removed, and the residue was purified by silica gel column chromatography to give 0.22 g of the objective compound.

Example 47

Production of (E)-N-methyl-2-[2'-((1"R*,2"R*)-1",2"-epoxy-2"-phenylethyl)phenyl]-2-methoxyiminoacetamide (Compound No. 116):-

In the same manner as in Example 45, the objective compound was prepared from N-methyl-2-(2-(E)-styrylphenyl)-2-(E)-methoxyiminoacetamide.


Example 48

Production of (E)-N-methyl-2-(2-phenylethylphenyl)-2-methoxyiminoacetamide (Compound No. 117):-

To a solution of N-methyl-2-(2-(E and Z)-styrylphenyl)-2-(E)-methoxyiminoacetamide or its isomer (0.30 g) in THF (5 ml), 10 % palladium-carbon (0.03 g) was added, and the resultant mixture was stirred at room temperature for 1 hour under hydrogen atmosphere. The reaction mixture was filtered, and the solvent was removed. The residue was purified by silica gel column chromatography to give 0.29 g of the objective compound.


In the same manner as above, Compounds Nos. 113 to 119 as shown in Table 10 below were obtained.

Table 10



Compound No.	Z	Steric configuration	m.p. (°C)	¹ H-NMR δ (CDCl ₃) (ppm)
113		E	165-166	2.94 (3H, d, J = 4.9), 3.95 (3H, s), 6.78 (1H, brs), 6.88 (1H, d, J = 15.6), 6.88 (1H, d, J = 15.6), 7.04 (1H, d, J = 15.6), 7.16-7.46 (8H, m), 7.74 (1H, d, J = 6.8)
114		E	94-95	2.89 (3H, d, J = 5.9), 3.84 (3H, s), 6.44 and 6.55 (each 1H, ABq, J = 11.7), 6.64 (1H, brs), 7.15-7.29 (9H, m)
115		E	oily	2.88 (3H, d, J = 4.9), 3.93 (3H, s), 4.22 (1H, d, J = 3.9), 4.29 (1H, d, J = 3.9), 6.54 (1H, brs), 7.04-7.23 (9H, m)

(Continued)

116		E	164-166	2.82 (3H, d, J = 5.1), 3.66 (1H, d, J = 2.0), 3.72 (3H, s), 3.76 (1H, d, J = 2.0), 6.67 (1H, brs), 7.20 (1H, d, J = 7.3), 7.30-7.46 (8H, m)
117	-CH ₂ CH ₂ -	E	130-131	2.70-2.87 (4H, m), 2.92 (3H, d, J = 5.1), 3.95 (3H, s), 6.72 (1H, brs), 7.08-7.33 (9H, m)
118	-CH(OH)-	E	102-103	2.89 (3H, brd), 3.92 (3H, brs), 5.67 (1H, brs), 7.06-7.39 (10H, m)
119	-CO-	E	oily	2.85 (3H, d, J = 4.9), 3.76 (3H, s), 6.75 (1H, brs), 7.42-7.81 (9H, m)

Some examples for production of methyl-2-(2-phenoxyphenyl)-2-oxo-acetate, which is the intermediate compound for methyl-2-(2-phenoxyphenyl)-2-methoxyiminoacetamide, are as follows:

Reference Example 1

(1) 85 % Potassium hydroxide powders (2.18 g; 0.033 mol, 1.1 equimolar amount) were added to phenol (4.23 g; 0.045 mol, 1.5 eq.) under heating, and 1,2-dichlorobenzene (4.41 g; 0.03 mol) and copper(I) iodide (0.29 g; 0.0015 mol) were added thereto, followed by stirring at 160°C for 24 hours. The reaction mixture was diluted with water and extracted with ether. The extract was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with a mixture of benzene and n-hexane to give 3.48 g (yield, 56.7 %) of 2-chlorodiphenyl ether as colorless crystals. m.p., 33 to 41°C.

(2) A mixture of magnesium (0.49 g; 0.02 mol) in dry THF (3 ml) and a slight amount of iodide was heated at 60°C for 5 minutes under argon stream, and a mixture of 2-chlorodiphenyl ether as above obtained (2.05 g; 0.01 mol) and dry THF (12 ml) was dropwise added thereto at 50 to 60°C in 10 minutes, followed by refluxing for 6 hours. The reaction mixture was dropwise added to a solution of dimethyl oxalate (1.18 g; 0.015 mol) in dry THF (15 ml) at not more than -3°C in 2 minutes, followed by stirring at room temperature overnight. An aqueous ammonium chloride solution was poured into the reaction mixture, which was extracted with ether. The extract was washed with water, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with a mixture of ethyl acetate and n-hexane to give 1.70 g (yield, 66.4 %) of the objective compound as a colorless oil.

Reference Example 2

(1) To a mixture of 2-phenoxybenzoic acid (7.71 g; 0.036 mol) dry benzene (80 ml), thionyl chloride (4.76 g; 0.04 mol) and DMF (2 drops) were added, and the resultant mixture was stirred under reflux for 2 hours. The reaction mixture was dropwise added to dry ethanol (100 ml) under ice-cooling in 30 minutes, followed by allowing to stand overnight. The reaction mixture was concentrated under reduced pressure and, after addition of a 4 % aqueous sodium hydrogen carbonate solution (150 ml) thereto, extracted with methylene chloride. The extract was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give ethyl-2-phenoxybenzoate (8.72 g, yield, 100 %) as a colorless oil.

(2) To a mixture of magnesium (2.19 g; 0.09 mol), dry benzene (36 ml) and dry THF (14.5 ml), methyl iodide (12.77 g; 0.09 mol) was added under ice-cooling for 2 minutes, followed by stirring at room temperature for 1 hour. To the resultant mixture, triethylamine (27.32 g; 0.27 mol) and a solution of ethyl 2-phenoxybenzoate as above obtained (8.72 g; 0.036 mol) in dry benzene (36 ml) were added at not more than 10°C for 40 minute, followed by stirring at 10°C for 4 hours. The reaction mixture was combined with ether (400 ml), washed with a 4 % aqueous hydrochloric acid solution (300 ml) and a 4 % aqueous sodium hydrogen carbonate solution (300 ml), dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography with a mixture of benzene and n-hexane to give 3.82 g (yield, 50.0 %) of 2-phenoxyacetophenone as a pale brown oil.

(3) To a solution of 2-phenoxyacetophenone obtained in (2) above (2.12 g; 0.01 mol) in dry methanol (2 ml) and dry chloroform (18 ml), bromide (1.60 g; 0.01 mol) was added at 40°C in 5 minutes, and the resultant mixture was stirred at room temperature for 30 minutes. The reaction mixture was combined with water (100 ml), extracted with methylene chloride, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give 3.02 g (yield, 102.3 %) of alpha-bromo-2-phenoxyacetophenone as a pale brown oil.

(4) The thus obtained alpha-bromo-2-phenoxyacetophenone (1.18 g; 0.004 mol) was added to a mixture of selenium dioxide (0.49 g; 0.0044 mol) and dry methanol (4 ml) under reflux, and the resultant mixture was stirred under reflux for 18 hours. Insoluble materials were removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography with a mixture of ethyl acetate and n-hexane to give 0.76 g (yield, 74.1 %) of methyl-2-(2-phenoxyphenyl)-2-oxo-acetate as a colorless oil.

Practical embodiments of the fungicidal composition according to the invention are illustratively shown in the following Formulation Examples wherein part(s) are by weight. The compound number as the active ingredient corresponds to the one in Tables 1 to 10 supra.

Preparation Example 1

Two parts of Compound No. 3 and 98 parts of talc are mixed well and pulverised to obtain powders.

Preparation Example 2

Forty parts of Compound No. 3, 10 parts of sodium lignin sulfonate and 50 parts of water are mixed well to obtain a dispersion.

Preparation Example 3

Ten parts of Compound No. 3, 1 part of Tween 20^R and 89 parts of isopropanol are mixed well to obtain a solution.

5 Preparation Example 4

Fifty parts of Compound No. 4, 6 parts of alkylbenzenesulfonate, 4 parts of sodium lignin sulfonate and 40 parts of clay are mixed well and pulverized to obtain a wettable powder.

10 Preparation Example 5

Five parts of Compound No. 3, 90 parts of an equilibrated mixture of bentonite and talc and 5 parts of alkylbenzene sulfonate are mixed well and pulverized to obtain granules.

15 Preparation Example 6

Twenty-five parts of Compound No. 1, 8 parts of polyoxyalkylphenyl ether, 2 parts of alkylbenzene sulfonate and 65 parts of xylene are dissolved to obtain an emulsifiable concentrate.

Typical biological data indicating the excellent fungicidal activity of the compounds of the invention are shown in the following Test Examples wherein a preparation containing methyl-2-phenoxy-phenyl-2-methoxy iminoacetate (JP-A-63-023852, EP-A-0254426) as the active compound is used for comparison, which is hereinafter referred to as "known compound".

(1) Controlling effect on plant diseases by foliar treatment (pot experiment)

25 Experiment 1

Controlling effect on Pyricularia oryzae:-

Two-week rice seedlings (var.: AICHISAH1) were transplanted in plastic cups (each 9 cmØ) and cultivated for another 2 weeks. The test compound in the form of a solution or a suspension was sprayed to the foliage of the rice seedlings, to which a conidia suspension of Pyricularia oryzae cultured in an oatmeal medium was inoculated by spraying. The test plant was kept in a moist chamber (28°C, 100% R.H.) for 24 hours, followed by cultivation in a greenhouse for 5 days. Six days after inoculation, the number of lesions for each plant was assessed and the percent control, i.e. preventive effect and curative effect, was calculated as follows:

Unless otherwise specifically mentioned, the preventive effect is determined by spraying a designated preparation the test plant and inoculating the pathogenic fungi thereto after 24 hours, followed by evaluation. The curative effect is determined by inoculating the pathogenic fungi to the test plants and spraying the designated preparation on the plant at the time of slight disease symptoms being observed, i.e. 24 to 48 hours after inoculation, followed by evaluation. The designated preparation herein is the one obtained by dissolving the active compound in a small amount of N,N-dimethylformamide and diluting with water containing a spreading agent to a desired concentration. The percent control was calculated on the following equation:

Percent control (%) =

$$\frac{\text{Severity or number of lesions in untreated plot} - \text{Severity or number of lesions in treated plot}}{\text{Severity or number of lesions in untreated plot}} \times 100$$

The results are shown in Table 11.

Table 11: Controlling effect on
Pyricularia oryzae

Compound No.	Percent control (%)		
	Preventive effect		Curative effect
	125 ppm	500 ppm	125 ppm
1	50	97	
3	97	97	97
4	50	90	90
5	30	97	50
6	97	100	97
7	50	90	50
8	90	97	97
16	90	90	90
17	70	90	70
19		70	
21		70	
22	70	90	90
29		70	
33	97	90	90
36	97	97	90
39	97	90	97
41	97	97	97
46	90	90	97
47	90	90	90
48	90	90	90
49	90	90	90
50	90	97	90
51	90	97	97
54	70	90	70
55	90	97	97
57	90	97	90
58	90	90	97
59	70	90	50
60	97	97	90
63		70	
64	50	97	97
65	90	90	97
66	70	97	97
67	70	90	70
71	30	97	90
72	50	97	70

(Continued)

5	73	70	97	70
	75	50	97	50
	82	70	90	97
	84	97	90	90
	85	50	90	70
10	86	90	90	97
	87	90	90	90
	88		70	
	97		70	
	98		70	
15	103		70	
	104	70	90	90
	105	70	97	97
	108		70	
	109		70	
20	110	97	97	97
	113	70	90	97
	Known compound		0	

Experiment 2Controlling effect on Rhizoctonia solani:-

Two-week rice seedlings (var.: AICHISAH) were transplanted in plastic cups (each 9 cmØ) and cultivated another 2 weeks. The test compound in the form of a solution or a suspension was sprayed to the sheath and foliage of the rice seedlings. Mycelia of Rhizoctonia solani, which were previously cultivated on the rice bran medium, were put at the feet of the seedlings, and the plant was kept in a moist chamber (28°C, 100% R.H.) for 5 days. The infectious state of the plant was assessed by measuring the height of the mycelia raised along the leaf sheath, and the percent control was calculated in the same manner as in Experiment 1. The results are shown in Table 12.

Table 12

Controlling effect on <u>Rhizoctonia solani</u>	
Compound No.	Preventive effect at 500 ppm (%)
3	70
4	50
6	90
7	70
8	70
16	50
19	70
22	70
41	70
46	90
Known compound	0

Experiment 3

Controlling effect on Sphaerotheca fuliginea:-

Seeds of cucumber (var.: TSUKUBASHIROIBO) were sown in plastic cups (each 9 cmØ), followed by cultivation for 2 to 3 weeks. The test compound in the form of a solution or suspension was sprayed on the surface of their first leaves, and a conidia suspension of Sphaerotheca fuliginea cultured on the cucumber leaves was sprayed thereto, and the plants were kept in a greenhouse at 20°C for 10 days. The controlling effect was assessed by observing the infected area on the leaves and calculated in the same manner as in Experiment 1. The results are shown in Table 13.

Table 13: Controlling effect on
Sphaerotheca fuliginea

Compound No.	Percent control (%)		
	Preventive effect		Curative effect
	125 ppm	500 ppm	125 ppm
3	100	100	100
4	100	100	97
5	100	100	100
6	100	100	100
7	100	100	100
8	100	100	100
11		70	
16	100	100	100
17	100	100	100
19	100	100	98
21		70	
22	100	100	100
28	90	97	95
29	100	100	100
30		70	
31	100	100	100
32	100	97	90
33	100	100	98
34	100	100	100
35	70	97	50
36	100	100	100
39	100	100	100
41	100	100	100
44		70	
46	97	100	100
47	100	100	100
48	100	100	100
49	100	100	100
50	100	100	
51	100	100	100
57	100	100	100
58	100	100	100
59	100	100	100
60	100	100	100
62	100	100	100
64	100	100	100
65	100	100	100
66	100	100	100

(Continued)

5	67	100	100	100
	71	100	100	100
	72	85	90	100
	73	100	100	100
	75	100	100	100
10	82	100	97	100
	84	100	100	100
	85	100	100	75
	86		70	
	87	100	100	100
15	88	100	100	100
	89		70	
	90	100	100	90
	97	100	100	100
20	103	70	100	80
	105	100	100	100
	107	80	97	80
	108	100	100	100
	109	100	100	100
25	110	100	100	100
	known compound	100	100	100

Experiment 4Controlling effect on *Pseudoperonospora cubensis*:-

The seeds of cucumber (var.: TSUKUBASHIROIBO) were sown in plastic cups (9 cmØ), followed by cultivation for 2 to 3 weeks. The test compound in the form of a solution or suspension was sprayed to the surface of their first leaves. A zoospore suspension of *Pseudoperonospora cubensis* cultured on the cucumber leaves was dropped on the under surface (untreated side) of the leaves, and the plants were cultivated in a greenhouse at 20°C for 10 days. The controlling effect was assessed by observing the infected area on the leaves and calculated in the same manner as in Experiment 1.

The results are shown in Table 14.

Table 14

Controlling effect on <u>Pseudoperonospora cubensis</u>	
Compound No.	Preventive effect at 500 ppm (%)
3	100
6	70
8	70
16	70
31	70
33	70
39	70
41	70
46	70
47	70
48	70
50	70
51	70
58	70
64	100
65	100
66	100
71	70
72	100
75	70
103	97
105	100
107	70
110	70
Known compound	0

Experiment 5Controlling effect on Botrytis cinerea:-

The seeds of cucumber (var.: TSUKUBASHIROIBO) were sown in plastic cups (9 cmØ), followed by cultivation for 2 to 3 weeks. The test compound in the form of a solution or suspension was sprayed on the surface of their first leaves, and the cucumber seedlings were inoculated with mycelial disks (4 mmØ) of Botrytis cinerea cultured on the potato sucrose agar medium by putting the disks on the leaf surfaces. The plants were cultivated in a moist chamber at 20°C for 2 days. The percent control was assessed by measuring the diameter of the lesions on the leaves and calculated in the same manner as in Experiment 1. The results are shown in Table 15.

Table 15: Controlling effect on Botrytis cinerea

Compound No.	Percent control (%)	
	Preventive effect	Curative effect
	500 ppm	500 ppm
3	96	94
5	50	93
6	74	95
7	71	93
8	68	93
9	50	74
11	70	89
15	59	94
16	100	94
17	100	95
19	74	90
20	51	82
21	70	92
22	90	92
29	100	98
31	100	94
32	100	93
33	100	95
34	100	98
35	50	
36	97	96
39	100	87
41	70	83
45	71	51
46	100	94
47	70	54
48	70	96
49	70	97

(Continued)

5	50	72	92
	51	86	94
	53	63	86
	55	79	96
	56	50	
	57	74	93
10	58	90	93
	59	70	
	60	70	95
	61	50	
	62	70	96
15	63	50	
	64	70	92
	65	70	94
	66	90	99
	67	70	94
20	71	70	95
	72	90	94
	73	90	98
	75	62	96
	82	50	
25	84	50	
	85	62	86
	86	58	81
	87	63	92
30	88	70	68
	89	50	
	90	69	71
	91	60	86
	92	59	77
35	94	50	
	95	70	62
	97	70	
	98	50	
	103	70	
40	104	70	
	105	100	
	107	50	
	108	66	90
	109	42	86
45	110	93	94
	113	86	94
	Known compound	100	65

50

The preparations comprising the compounds according to the invention showed in all tested diseases the controlling effect as equal to, or remarkably higher than the known compound, so that their fungicidal activity was confirmed to be higher and their fungicidal spectrum was broader than those of the known compound.

55

Various compounds, which have been conventionally used as agricultural fungicides such as Captan and Dithane having a broad fungicidal spectrum but not systemic, are useful. However, their fungicidal activity is only exerted on treatment before infection, and any systemic activity during the growth of plants is hardly expected and are thus called the protective fungicides. On the other hand, the compounds of the invention not only have a broad fungicidal spectrum over the wide range of the pathogenic fungi but also possess a high systemic activity. Thus, the compounds according

to the invention are useful in protecting the agricultural plants from the attack of the pathogenic fungi with remarkable preventive and curative effects.

(2) Controlling effect on *Pyricularia oryzae* by paddy water application (pot experiment)

Experiment 6

Preventive effect on *Pyricularia oryzae*:-

Seeds of rice plant (var.: AICHIASAHI) were sowed in plastic cups (9 cmØ) and cultivated in a greenhouse at 28°C for 9 days. The test compound dissolved in a small amount of acetone gas dropped into the flooded cups at a concentration of the active ingredient being 100 ppm at the maximum. The rice seedlings treated thus were further grown in a greenhouse for 7 days. A conidia suspension of *Pyricularia oryzae* cultivated in an oatmeal medium was sprayed on the foliage of the plants, which were kept in a moist chamber (28°C, 100% R.H.) for 24 hours, followed by cultivation in the greenhouse for 4 days. The number of lesions were observed and the percent control was calculated in the same manner as in Experiment 1. The results are shown in Table 16.

Table 16


Controlling effect on <i>Pyricularia oryzae</i>	
Compound No.	Percent control (%) at 10 ppm
1	70
3	97
4	97
5	97
6	100
7	99
8	100
16	97
17	97
55	98
82	99
Known compound	0

It is obvious from the above test results that the compounds of this invention show a remarkable controlling effect by paddy water application in comparison with the known compound. This prominent fungicidal activity is attributable to the uptake of the compound from the root of the plant.


Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE,

1. A compound of the formula:




wherein R^1 and R^2 are each independently hydrogen, C_1 - C_8 alkyl or cyclo(C_3 - C_8)alkyl; R^3 is C_1 - C_8 alkyl or cyclo(C_3 - C_8)alkyl; R^4 and R^5 are each hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen-substituted C_1 - C_8 alkyl, C_1 - C_8 alkyl-substituted silyl, halogen or nitro; A represents an unsaturated hydrocarbon group, a halogen-substituted unsaturated hydrocarbon group, a phenyl group or a heterocyclic group, said phenyl group or heterocyclic group being optionally substituted with not more than three substituents; and Z is $-CH_2-$, $-CH(OH)-$, $-CO-$, $-O-$, $-S-$, $-SO-$, $-NR-$ (R being hydrogen or C_1 - C_8 alkyl), $-CH_2CH_2-$, $-CH=CH-$,




$-CH_2O-$, $-CH_2S-$, $-CH_2SO-$, $-OCH_2-$, $-SCH_2-$ or $-SOCH_2-$, provided that when Z is O A is other than 2-pyridyl.

2. A compound of the formula:




wherein Z is $-O-$, $-CH_2O-$, $-OCH_2-$, $-S-$, $-SO-$, $-SCH_2-$ or $-SOCH_2-$, R^1 and R^2 are each independently hydrogen or C_1 - C_8 alkyl, R^3 is C_1 - C_8 alkyl, R^4 and R^5 are each independently hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen-substituted C_1 - C_8 alkyl, C_1 - C_8 alkyl-substituted silyl, halogen or nitro and X_1 , X_2 and X_3 are each hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, C_1 - C_8 alkanoyl, halogen-substituted C_1 - C_8 alkyl, C_1 - C_8 alkyl-substituted silyl, halogen, nitro, cyano, di(C_1 - C_8)alkylamino, phenyl, phenyl(C_2 - C_8)alkenyl, furyl(C_2 - C_8)alkenyl, hydroxyl, C_2 - C_8 alkynyloxy, C_1 - C_8 alkanoyloxy, benzoyloxy, phenoxy, C_1 - C_8 alkoxyphenoxy, nitrophenoxy, benzyloxy, cyanobenzyloxy, tetrahydropyran-
nyloxy, pyrimidinyl, pyridyl, trifluoromethylpyridyl, benzothiazolyl, quinolyl, benzoyl(C_1 - C_8)alkoxy, benzenesulfonyloxy or C_1 - C_8 alkylbenzenesulfonyloxy.

3. A compound as claimed in claim 2, wherein Z is oxygen, R¹ is hydrogen, R² is C₁-C₆ alkyl, R³ is C₁-C₆ alkyl and R⁴, R⁵, X₁, X₂ and X₃ are each as defined in claim 2.
4. A compound as claimed in claim 1, wherein Z is oxygen, R¹ is hydrogen, R² is C₁-C₆ alkyl, R³ is C₁-C₆ alkyl, R⁴ is hydrogen, R⁵ is hydrogen and A is phenyl.
5. A compound according to claim 1, wherein Z is O, R¹ is hydrogen, R² is methyl, R³ is methyl, R⁴ is hydrogen, R⁵ is hydrogen and A is phenyl.
6. A compound as claimed in claim 1, which has the formula:




wherein R¹ and R² are each hydrogen or C₁-C₆ alkyl, R³ is C₁-C₆ alkyl, R⁴ and R⁵ are each hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen-substituted C₁-C₆ alkyl, C₁-C₆ alkyl-substituted silyl, halogen or nitro, R⁷ is phenyl, C₁-C₆ alkoxyphenyl, halophenyl, pyridyl or pyrimidinyl and Z' is -O-, -S-, -SO- or -NR- (R being C₁-C₆ alkyl).

7. A compound as claimed in claim 1, which has the formula:



wherein R¹ and R² are each hydrogen or C₁-C₆ alkyl, R³ is C₁-C₆ alkyl, A' is pyridyl or pyrimidinyl, Z is -O-, -OCH₂- or -CH₂O- and X₁ is hydrogen or trifluoromethyl.

8. A compound as claimed in claim 1, which has the formula:




15 wherein R¹ and R² are each hydrogen or C₁-C₆ alkyl, R³ is C₁-C₆ alkyl and Z is -CH₂CH₂-,



20 -CH=CH-, -CH(OH)- or -CO-.

9. A compound as claimed in claim 1, which has the formula:



35 wherein R¹ and R² are each hydrogen or C₁-C₆ alkyl, R³ is C₁-C₆ alkyl and A is alkenyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms or alkadienyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms.


40 10. A process for preparing a compound as claimed in any one of claims 1 to 9, which comprises subjecting a compound of the formula:



55 wherein R⁴, R⁵, A and Z are each as defined in claim 1 to amidation and alkoximation, or vice versa.

11. A process as claimed in claim 10, wherein amidation is carried out first and then alkoximation is effected.


12. A process as claimed in claim 10, wherein alkoxylation is carried out first and then amidation is effected.
13. A process as claimed in any one of claims 10 to 12 wherein the resulting compound is isolated and admixed with one or more components selected from carriers, diluents, surfactants, adherents, dispersants and stabilizers.
14. A fungicidal composition or formulation which comprises as an active ingredient at least one compound as claimed in any one of claims 1 to 9, together with one or more components selected from carriers, diluents, surfactants, adherents, dispersants and stabilizers, which said component may be inert, optionally further comprising at least one other fungicide, insecticide, herbicide or fertilizer.
15. A method for controlling phytopathogenic fungi which comprises applying as an active ingredient at least one of the compounds according to any one of claims 1 to 9 to a locus where phytopathogenic fungi may propagate or are propagating.
16. The use of a compound as claimed in any one of claims 1 to 9 as a fungicide.
17. The use of a compound of the formula:



wherein R^1 and R^2 are each independently hydrogen, C_1 - C_8 alkyl or cyclo (C_3 - C_8) alkyl; R^3 is C_1 - C_8 alkyl or cyclo (C_3 - C_8) alkyl; R^4 and R^5 are each hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen-substituted C_1 - C_8 alkyl, C_1 - C_8 alkyl-substituted silyl, halogen or nitro; A represents pyridyl being optionally substituted with not more than three substituents and Z is -O- as a fungicide.

18. A compound as claimed in claim 3, wherein R^2 is methyl, R^3 is methyl, X_1 is hydrogen, X_2 is hydrogen and X_3 is either hydrogen or 4-methyl.
19. A compound as claimed in claim 2, wherein Z is $-OCH_2-$, R^1 is hydrogen and R^2 is methyl, R^3 is methyl, R^4 and R^5 are both hydrogen, and X_1 is hydrogen, and X_2 and X_3 are hydrogen, methyl, chlorine or fluoromethyl.
20. A compound as claimed in claim 19, wherein X_2 and X_3 are hydrogen or methyl.
21. A compound as claimed in claim 19, wherein X_2 is hydrogen and X_3 is 3- CF_3 , or X_2 is hydrogen and X_3 is 2-Cl or X_2 is hydrogen and X_3 is 4-Cl.


22. A compound of formula:




15 wherein X is O or N-OH, and A, Z, R¹, R², R³, R⁴ and R⁵ are as defined in claim 1.

Claims for the following Contracting States : ES, GR


20 1. A process for preparing a compound of the formula:



35 wherein R¹ and R² are each independently hydrogen, C₁-C₈ alkyl or cyclo(C₃-C₈)alkyl; R³ is C₁-C₈ alkyl or cyclo (C₃-C₈) alkyl; R⁴ and R⁵ are each hydrogen, C₁-C₈ alkyl, C₁-C₈ alkoxy, halogen-substituted C₁-C₈ alkyl, C₁-C₈ alkyl-substituted silyl, halogen or nitro; A represents an unsaturated hydrocarbon group, a halogen-substituted unsaturated hydrocarbon group, a phenyl group or a heterocyclic group, said phenyl group or heterocyclic group being optionally substituted with not more than three substituents; and Z is -CH₂-, -CH(OH)-, -CO-, -O-, -S-, -SO-, -NR- (R being hydrogen or C₁-C₈ alkyl), -CH₂CH₂-, -CH=CH-,




50 -CH₂O-, -CH₂S-, CH₂SO-, -OCH₂-, -SCH₂- or -SOCH₂-, provided that when Z is O A is other than 2-pyridyl which process comprises subjecting a compound of the formula:



wherein R^4 , R^5 , A and Z are each as defined in claim 1 to amidation and alkoximation, or vice versa.


- 15
2. A process as claimed in claim 1, wherein amidation is carried out first and then alkoximation is effected.
 3. A process as claimed in claim 1, wherein alkoximation is carried out first and then amidation is effected.
 - 20 4. A process as claimed in any one of the preceding claims wherein the reactants are chosen to provide a compound of the formula:



wherein Z is -O-, -CH₂O-, -OCH₂-, -S-, -SO-, -SCH₂- or -SOCH₂-, R^1 and R^2 are each independently hydrogen or C₁-C₈ alkyl, R^3 is lower alkyl, R^4 and R^5 are each independently hydrogen, C₁-C₈ alkyl, C₁-C₈ alkoxy, halogen-substituted C₁-C₈ alkyl, C₁-C₈ alkyl-substituted silyl, halogen or nitro and X_1 , X_2 and X_3 are each hydrogen, C₁-C₈ alkyl, C₁-C₈ alkoxy, C₁-C₈ alkanoyl, halogen-substituted C₁-C₈ alkyl, C₁-C₈ alkyl-substituted silyl, halogen, nitro, cyano, di(C₁-C₈)alkylamino, phenyl, phenyl(C₂-C₈)alkenyl, furyl(C₂-C₈)alkenyl, hydroxyl, C₂-C₈ alkynyloxy, C₁-C₈ alkanoyloxy, benzoyloxy, phenoxy, C₁-C₈ alkoxyphenoxy, nitrophenoxy, benzyloxy, cyanobenzyloxy, tetrahydropyran-4-yloxy, pyrimidin-2-yloxy, pyridin-2-yloxy, trifluoromethylpyridin-2-yloxy, benzothiazol-2-yloxy, quinolin-2-yloxy, benzoyl(C₁-C₈)alkoxy, benzenesulfonyloxy or lower alkylbenzenesulfonyloxy.


- 45
5. A process as claimed in claim 4, wherein Z is oxygen, R^1 is hydrogen, R^2 is C₁-C₆ alkyl, R^3 is C₁-C₆ alkyl and R^4 , R^5 , X_1 , X_2 and X_3 are each as defined in claim 2.
 6. A process as claimed in claim 1, wherein Z is oxygen, R^1 is hydrogen, R^2 is C₁-C₆ alkyl, R^3 is C₁-C₆ alkyl, R^4 is hydrogen, R^5 is hydrogen and A is phenyl.
 7. A process according to claim 1, wherein Z is O, R^1 is hydrogen, R^2 is methyl, R^3 is methyl, R^4 is hydrogen, R^5 is hydrogen and A is phenyl.
- 55

8. A process as claimed in claim 1, wherein the reactants are chosen to provide a compound which has the formula:




wherein R^1 and R^2 are each hydrogen or C_1 - C_6 alkyl, R^3 is C_1 - C_6 alkyl, R^4 and R^5 are each hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen-substituted C_1 - C_6 alkyl, C_1 - C_6 alkyl-substituted silyl, halogen or nitro, R^7 is phenyl, C_1 - C_6 alkoxyphenyl, halophenyl, pyridyl or pyrimidinyl and Z' is -O-, -S-, -SO- or -NR- (R being C_1 - C_6 alkyl).

9. A process as claimed in claim 1, wherein the reactants are chosen to provide a compound which has the formula:




wherein R^1 and R^2 are each hydrogen or C_1 - C_6 alkyl, R^3 is C_1 - C_6 alkyl, A' is pyridyl or pyrimidinyl, Z is -O-, -OCH₂- or -CH₂O- and X_1 is hydrogen or trifluoromethyl.

10. A process as claimed in claim 1, wherein the reactants are chosen to provide a compound which has the formula:




wherein R^1 and R^2 are each hydrogen or C_1 - C_6 alkyl, R^3 is C_1 - C_6 alkyl and Z is -CH₂CH₂-,



-CH=CH-, -CH(OH)- or -CO-.

11. A process as claimed in claim 1, wherein the reactants are chosen to provide a compound which has the formula:



15 wherein R¹ and R² are each hydrogen or C₁-C₆ alkyl, R³ is C₁-C₆ alkyl and A is alkenyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms or alkadienyl of 3 to 10 carbon atoms optionally substituted with not more than 3 halogen atoms.

20 12. A process as claimed in any one of the preceding claims, wherein the resulting compound is isolated and admixed with one or more components selected from carriers, diluents, surfactants, adherents, dispersants and stabilizers.


25 13. A process for producing a fungicidal composition or formulation which comprises mixing an active ingredient at least one compound as defined in any one of claims 1 to 11, together with one or more components selected from carriers, diluents, surfactants, adherents, dispersants and stabilizers, which said component may be inert, optionally further comprising at least one other fungicide, insecticide, herbicide or fertilizer.

30 14. A method for controlling phytopathogenic fungi which comprises producing an active ingredient by carrying out a process as claimed in any one of claims 1 to 12, and thereafter applying the active ingredient to a locus where phytopathogenic fungi may propagate or are propagating.


Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE,

35 1. Verbindung mit der Formel,




55 wobei R¹ und R² jeweils unabhängig ein Wasserstoffatom, einen C₁-C₈-Alkyl- oder Cyclo(C₃-C₈)alkylrest darstellen, R³ ein C₁-C₈-Alkyl- oder Cyclo(C₃-C₈)alkylrest ist, R⁴ und R⁵ jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe bedeuten, A einen ungesättigten Kohlenwasserstoffrest, einen halogensubstituierten ungesättigten Kohlenwasserstoffrest, eine Phenylgruppe oder einen heterocyclischen Rest darstellt, wobei die Phenylgruppe oder der heterocyclische Rest gegebenenfalls mit nicht mehr als drei Substituenten substituiert ist, und Z -CH₂-, -CH(OH)-, -CO-, -O-, -S-, -SO-, -NR- (wobei R ein Wasserstoffatom oder ein C₁-C₈-Alkylrest ist), -CH₂CH₂-, -CH=CH-,



-CH₂O-, -CH₂S-, -CH₂SO-, -OCH₂-, -SCH₂- oder -SOCH₂- ist, mit der Maßgabe, daß, wenn Z ein Sauerstoffatom ist, A keine 2-Pyridylgruppe ist.

2. Verbindung mit der Formel




wobei Z -O-, -CH₂O-, -OCH₂-, -S-, -SO-, -SCH₂- oder -SOCH₂- ist, R¹ und R² jeweils unabhängig ein Wasserstoffatom oder einen C₁-C₈-Alkylrest darstellen, R³ ein C₁-C₈-Alkylrest ist, R⁴ und R⁵ jeweils unabhängig ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe darstellen und X₁, X₂ und X₃ jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, C₁-C₈-Alkanoyl-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom, eine Nitrogruppe, eine Cyanogruppe, einen Di(C₁-C₈)alkylaminorest, eine Phenylgruppe, einen Phenyl(C₂-C₈)alkenyl-, Furyl(C₂-C₈)alkenylrest, eine Hydroxylgruppe, einen C₂-C₈-Alkinyloxy-, C₁-C₈-Alkanoyloxyrest, eine Benzoyloxy-, Phenoxygruppe, einen C₁-C₈-Alkoxyphenoxyrest, eine Nitrophenoxy-, Benzyloxy-, Cyanobenzyloxy-, Tetrahydropyranyloxy-, Pyrimidinyl-, Pyridyl-, Trifluormethylpyridyl-, Benzothiazolyl-, Chinolyl-, einen Benzoyl(C₁-C₈)alkoxyrest, eine Benzolsulfonyloxygruppe oder einen C₁-C₈-Alkylbenzolsulfonyloxyrest darstellen.

3. Verbindung nach Anspruch 2, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² ein C₁-C₆-Alkylrest, R³ ein C₁-C₆-Alkylrest ist und R⁴, R⁵, X₁, X₂ und X₃ jeweils wie in Anspruch 2 definiert sind.

4. Verbindung nach Anspruch 1, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² ein C₁-C₆-Alkylrest, R³ ein C₁-C₆-Alkylrest, R⁴ ein Wasserstoffatom, R⁵ ein Wasserstoffatom und A eine Phenylgruppe ist.


5. Verbindung nach Anspruch 1, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² eine Methylgruppe, R³ eine Methylgruppe, R⁴ ein Wasserstoffatom, R⁵ ein Wasserstoffatom und A eine Phenylgruppe ist.

6. Verbindung nach Anspruch 1 mit der Formel




20 wobei R^1 und R^2 jeweils ein Wasserstoffatom oder einen C_1 - C_6 -Alkylrest darstellen, R^3 ein C_1 - C_6 -Alkylrest ist, R^4 und R^5 jeweils ein Wasserstoffatom, einen C_1 - C_6 -Alkyl-, C_1 - C_6 -Alkoxy-, halogensubstituierten C_1 - C_6 -Alkyl-, C_1 - C_6 -alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe darstellen, R^7 eine Phenylgruppe, einen C_1 - C_6 -Alkoxyphenyl-, Halogenphenylrest, eine Pyridyl- oder Pyrimidinylgruppe darstellt und Z' -O-, -S-, -SO- oder -NR- (wobei R ein C_1 - C_6 -Alkylrest ist), darstellt.

7. Verbindung nach Anspruch 1 mit der Formel




40 wobei R^1 und R^2 jeweils ein Wasserstoffatom oder ein C_1 - C_6 -Alkylrest sind, R^3 ein C_1 - C_6 -Alkylrest ist, A' eine Pyridyl- oder Pyrimidinylgruppe ist, Z -O-, -OCH₂- oder -CH₂O- ist und X_1 ein Wasserstoffatom oder eine Trifluormethylgruppe ist.

8. Verbindung nach Anspruch 1 mit der Formel




wobei R^1 und R^2 jeweils ein Wasserstoffatom oder einen C_1 - C_6 -Alkylrest darstellen, R^3 ein C_1 - C_6 -Alkylrest ist und Z -CH₂CH₂-,




-CH=CH-, -CH(OH)- oder -CO- ist.

9. Verbindung nach Anspruch 1 mit der Formel



wobei R¹ und R² jeweils ein Wasserstoffatom oder ein C₁-C₆-Alkylrest sind, R³ ein C₁-C₆-Alkylrest ist und A ein Alkenylrest mit 3 bis 10 Kohlenstoffatomen, gegebenenfalls substituiert mit nicht mehr als 3 Halogenatomen, oder ein Alkadienylrest mit 3 bis 10 Kohlenstoffatomen, gegebenenfalls substituiert mit nicht mehr als 3 Halogenatomen, ist.

10. Verfahren zur Herstellung einer Verbindung nach einem der Ansprüche 1 bis 9, umfassend die Amidierung und Alkoximierung, oder umgekehrt, einer Verbindung mit der Formel



wobei R⁴, R⁵, A und Z jeweils wie in Anspruch 1 definiert sind.

11. Verfahren nach Anspruch 10, wobei die Amidierung zuerst durchgeführt und die Alkoximierung danach vollzogen wird.

12. Verfahren nach Anspruch 10, wobei die Alkoximierung zuerst durchgeführt und die Amidierung danach vollzogen wird.

13. Verfahren nach einem der Ansprüche 10 bis 12, wobei die resultierende Verbindung isoliert und mit einer oder mehreren Komponenten vermischt wird, die aus Trägerstoffen, Verdünnungsmitteln, oberflächenaktiven Substanzen, Adhärenzen, Dispergierungsmitteln und Stabilisatoren gewählt werden.


14. Fungizide Zusammensetzung oder Zubereitung, welche als Wirkstoff mindestens eine Verbindung nach einem der Ansprüche 1 bis 9 umfaßt, zusammen mit einer oder mehreren Komponenten gewählt aus Trägerstoffen, Verdünnungsmitteln, oberflächenaktiven Substanzen, Adhärenzen, Dispergierungsmitteln und Stabilisatoren, wobei die

Komponente inert sein kann, und gegebenenfalls zusätzlich mindestens ein anderes Fungizid, Insektizid, Herbizid oder Düngemittel enthält.

15. Verfahren zur Bekämpfung phytopathogener Pilze, umfassend die Anwendung mindestens einer Verbindung nach einem der Ansprüche 1 bis 9 als wirksamen Bestandteil an einem Ort, wo sich phytopathogene Pilze vermehren können oder sich vermehren.

16. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 9 als Fungizid.

17. Verwendung einer Verbindung mit der Formel



wobei R^1 und R^2 jeweils unabhängig ein Wasserstoffatom, einen C_1 - C_8 -Alkyl- oder Cyclo(C_3 - C_8)alkylrest darstellen, R^3 ein C_1 - C_8 -Alkyl- oder Cyclo(C_3 - C_8)alkylrest ist, R^4 und R^5 jeweils ein Wasserstoffatom, einen C_1 - C_8 -Alkyl-, C_1 - C_8 -Alkoxy-, halogensubstituierten C_1 - C_8 -Alkyl-, C_1 - C_8 -alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe bedeuten, A eine Pyridylgruppe darstellt, welche gegebenenfalls mit nicht mehr als drei Substituenten substituiert ist und Z ein Sauerstoffatom ist, als Fungizid.


18. Verbindung nach Anspruch 3, wobei R^2 eine Methylgruppe, R^3 eine Methylgruppe, X_1 ein Wasserstoffatom, X_2 ein Wasserstoffatom und X_3 entweder ein Wasserstoffatom oder eine 4-Methylgruppe ist.

19. Verbindung nach Anspruch 2, wobei Z $-OCH_2-$ ist, R^1 ein Wasserstoffatom und R^2 eine Methylgruppe, R^3 eine Methylgruppe ist, R^4 und R^5 beide ein Wasserstoffatom sind und X_1 ein Wasserstoffatom ist und X_2 und X_3 ein Wasserstoffatom, eine Methylgruppe, ein Chloratom oder eine Fluormethylgruppe bedeuten.

20. Verbindung nach Anspruch 19, wobei X_2 und X_3 ein Wasserstoffatom oder eine Methylgruppe darstellen.

21. Verbindung nach Anspruch 19, wobei X_2 ein Wasserstoffatom und X_3 eine 3- CF_3 -Gruppe ist, oder X_2 ein Wasserstoffatom und X_3 ein 2-Chloratom ist oder X_2 ein Wasserstoffatom und X_3 ein 4-Chloratom ist.


22. Verbindung mit der Formel




wobei X ein Sauerstoffatom oder eine N-OH-Gruppe ist und A, Z, R¹, R², R³, R⁴ und R⁵ wie in Anspruch 1 definiert sind.

Patentansprüche für folgende Vertragsstaaten : ES, GR


- Verfahren zur Herstellung einer Verbindung mit der Formel,



wobei R¹ und R² jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl- oder Cyclo(C₃-C₈)alkylrest darstellen, R³ ein C₁-C₈-Alkyl- oder Cyclo(C₃-C₈)alkylrest ist, R⁴ und R⁵ jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe bedeuten, A einen ungesättigten Kohlenwasserstoffrest, einen halogensubstituierten ungesättigten Kohlenwasserstoffrest, eine Phenylgruppe oder einen heterocyclischen Rest darstellt, wobei die Phenylgruppe oder der heterocyclische Rest gegebenenfalls mit nicht mehr als drei Substituenten substituiert ist, und Z -CH₂-, -CH(OH)-, -CO-, -O-, -S-, -SO-, -NR- (wobei R ein Wasserstoffatom oder ein C₁-C₈-Alkylrest ist), -CH₂CH₂-, -CH=CH-,




-CH₂O-, -CH₂S-, -CH₂SO-, -OCH₂-, -SCH₂- oder -SOCH₂- ist, mit der Maßgabe, daß, wenn Z ein Sauerstoffatom ist, A keine 2-Pyridylgruppe ist, umfassend die Amidierung und Alkoximierung, oder umgekehrt, einer Verbindung mit der Formel:



wobei R⁴, R⁵, A und Z jeweils wie in Anspruch 1 definiert sind.


- Verfahren nach Anspruch 1, wobei die Amidierung zuerst durchgeführt und die Alkoximierung danach vollzogen wird.
- Verfahren nach Anspruch 1, wobei die Alkoximierung zuerst durchgeführt und die Amidierung danach vollzogen wird.

4. Verfahren nach einem der Ansprüche 1 bis 3, wobei die resultierende Verbindung die Formel hat:




worin Z -O-, -CH₂O-, -OCH₂-, -S-, -SO-, -SCH₂- oder -SOCH₂- ist, R¹ und R² jeweils ein Wasserstoffatom oder einen C₁-C₈-Alkylrest darstellen, R³ ein Niederalkylrest ist, R⁴ und R⁵ jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe darstellen und X₁, X₂ und X₃ jeweils ein Wasserstoffatom, einen C₁-C₈-Alkyl-, C₁-C₈-Alkoxy-, C₁-C₈-Alkanoyl-, halogensubstituierten C₁-C₈-Alkyl-, C₁-C₈-alkylsubstituierten Silylrest, ein Halogenatom, eine Nitrogruppe, eine Cyanogruppe, einen Di(C₁-C₈)alkylaminorest, eine Phenylgruppe, einen Phenyl-(C₂-C₈)alkenyl-, Furyl-(C₂-C₈)alkenylrest, eine Hydroxylgruppe, einen C₂-C₈-Alkinyloxy-, C₁-C₈-Alkanoyloxyrest, eine Benzoyloxy-, Phenoxygruppe, einen C₁-C₈-Alkoxyphenoxyrest, eine Nitrophenoxy-, Benzyloxy-, Cyanobenzyloxy-, Tetrahydropyranlyoxy-, Pyrimidinlyoxy-, Pyridyloxy-, Trifluormethylpyridyloxy-, Benzothiazolyloxy-, Chinolyoxygruppe, einen Benzoyl(C₁-C₈)alkoxyrest, eine Benzolsulfonyloxygruppe oder einen Niederalkylbenzolsulfonyloxyrest darstellen.

5. Verfahren nach Anspruch 4, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² ein C₁-C₆-Alkylrest, R³ ein C₁-C₆-Alkylrest ist und R⁴, R⁵, X₁, X₂ und X₃ jeweils wie in Anspruch 2 definiert sind.
6. Verfahren nach einem der Ansprüche 1 bis 3, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² ein C₁-C₆-Alkylrest, R³ ein C₁-C₆-Alkylrest, R⁴ ein Wasserstoffatom, R⁵ ein Wasserstoffatom und A eine Phenylgruppe ist.
7. Verfahren nach einem der Ansprüche 1 bis 3, wobei Z ein Sauerstoffatom, R¹ ein Wasserstoffatom, R² eine Methylgruppe, R³ eine Methylgruppe, R⁴ ein Wasserstoffatom, R⁵ ein Wasserstoffatom und A eine Phenylgruppe ist.
8. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Ausgangsstoffe so gewählt werden, daß die resultierende Verbindung die Formel hat




wobei R¹ und R² jeweils ein Wasserstoffatom oder einen C₁-C₆-Alkylrest darstellen, R³ ein C₁-C₆-Alkylrest ist, R⁴ und R⁵ jeweils ein Wasserstoffatom, einen C₁-C₆-Alkyl-, C₁-C₆-Alkoxy-, halogensubstituierten C₁-C₆-Alkyl-, C₁-C₆-alkylsubstituierten Silylrest, ein Halogenatom oder eine Nitrogruppe darstellen, R⁷ eine Phenylgruppe, einen C₁-C₆-Alkoxyphenyl-, Halogenphenylrest, eine Pyridyl- oder Pyrimidinylgruppe darstellt und Z' -O-, -S-, -SO- oder -NR- (wobei R ein C₁-C₆-Alkylrest ist), darstellt.

9. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Ausgangsstoffe so gewählt werden, daß die resultierende Verbindung die Formel hat




15 wobei R¹ und R² jeweils ein Wasserstoffatom oder ein C₁-C₆-Alkylrest sind, R³ ein C₁-C₆-Alkylrest ist, A' eine Pyridyl- oder Pyrimidinylgruppe ist, Z -O-, -OCH₂- oder -CH₂O- ist und X₁ ein Wasserstoffatom oder eine Trifluormethylgruppe ist.

- 20 10. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Ausgangsstoffe so gewählt werden, daß die resultierende Verbindung die Formel hat




35 wobei R¹ und R² jeweils ein Wasserstoffatom oder einen C₁-C₆-Alkylrest darstellen, R³ ein C₁-C₆-Alkylrest ist und Z -CH₂CH₂-,



-CH=CH-, -CH(OH)- oder -CO-ist.

- 45 11. Verfahren nach einem der Ansprüche 1 bis 3, wobei die Ausgangsstoffe so gewählt werden, daß die resultierende Verbindung die Formel hat



wobei R¹ und R² jeweils ein Wasserstoffatom oder ein C₁-C₆-Alkylrest sind, R³ ein C₁-C₆-Alkylrest ist und A ein Alkenylrest mit 3 bis 10 Kohlenstoffatomen, gegebenenfalls substituiert mit nicht mehr als 3 Halogenatomen, oder


ein Alkadienylrest mit 3 bis 10 Kohlenstoffatomen, gegebenenfalls substituiert mit nicht mehr als 3 Halogenatomen ist.

12. Verfahren nach einem der Ansprüche 1 bis 11, wobei die resultierende Verbindung isoliert und dann mit einer oder mehreren Komponenten vermischt wird, die aus Trägerstoffen, Verdünnungsmitteln, oberflächenaktiven Substanzen, Adhärenzen, Dispergierungsmitteln und Stabilisatoren gewählt werden.
13. Verfahren zur Herstellung einer fungiziden Zusammensetzung, welches das Vermischen von mindestens einer resultierenden Verbindung nach einem der Ansprüche 1 bis 11 mit einer oder mehreren Komponenten gewählt aus Trägerstoffen, Verdünnungsmitteln, oberflächenaktiven Substanzen, Adhärenzen, Dispergierungsmitteln und Stabilisatoren umfaßt, wobei die Komponente inert sein kann, und gegebenenfalls zusätzlich mindestens ein anderes Fungizid, Insektizid, Herbizid oder Düngemittel enthält.
14. Verfahren zur Bekämpfung phytopathogener Pilze, umfassend die Herstellung eines Wirkstoffes durch Ausführung eines Verfahrens nach einem der Ansprüche 1 bis 12 und danach Anwendung des Wirkstoffes an einem Ort, wo sich phytopathogene Pilze vermehren können oder sich vermehren.


Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

1. Composé de la formule :




dans laquelle R^1 et R^2 représentent chacun indépendamment un atome d'hydrogène, un radical alkyle en C_1 à C_8 ou cycloalkyle en C_3 à C_8 , R^3 représente un radical alkyle en C_1 à C_8 ou cycloalkyle en C_3 à C_8 , R^4 et R^5 représentent chacun un atome d'hydrogène, un radical alkyle en C_1 à C_8 , un radical alcoxy en C_1 à C_8 , un radical alkyle en C_1 à C_8 à substitution halogénée, un radical silyle à substitution alkylque en C_1 à C_8 , un atome d'halogène, ou le radical nitro, A représente un groupe hydrocarboné insaturé, un radical hydrocarboné insaturé à substitution halogénée, un radical phényle, ou un radical hétérocyclique, le radical phényle précité ou le radical hétérocyclique précité étant éventuellement substitué par pas plus de trois substituants et Z représente un radical $-CH_2-$, $-CH(OH)-$, $-CO-$, $-O-$, $-S-$, $-SO-$, $-NR-$ (R représentant un atome d'hydrogène ou un radical alkyle en C_1-C_8), $-CH_2CH_2-$, $-CH=CH-$,




$-CH_2O-$, $-CH_2S-$, $-CH_2SO-$, $-OCH_2-$, $-SCH_2-$ ou $-SOCH_2-$ avec la condition que lorsque Z représente O, A soit autre que le radical 2-pyridyle.

2. Composé de la formule :



dans laquelle Z représente -O-, -CH₂O-, -OCH₂-, -S-, -SO-, -SCH₂- ou -SOCH₂-, R¹ et R² représentent chacun indépendamment un atome d'hydrogène ou un radical alkyle en C₁ à C₈, R³ représente un radical alkyle en C₁ à C₈, R⁴ et R⁵ représentent chacun indépendamment un atome d'hydrogène, un radical alkyle en C₁ à C₈, alcoxy en C₁ à C₈, un radical alkyle en C₁ à C₈ halogénosubstitué, un radical silyle à substitution alkylque en C₁ à C₈, un atome d'halogène, ou le radical nitro et X₁, X₂ et X₃ représentent chacun un atome d'hydrogène, un radical alkyle en C₁ à C₈, alcoxy en C₁ à C₈, alcanoyloxy en C₁ à C₈, alkyle en C₁ à C₈ à substitution halogénée, silyle à substitution alkylque en C₁ à C₈, un atome d'halogène, le radical nitro, cyano, un radical dialkyl(C₁ à C₈)phényle, phénylalcényloxy(C₂ à C₈), furylalcényloxy(C₂ à C₈), hydroxyle, alcynyloxy en C₂ à C₈, alcanoyloxy en C₁ à C₈, benzoyloxy, phénoxy, alcoxy(C₁ à C₈)phénoxy, nitrophénoxy, benzyloxy, cyanobenzyloxy, tétrahydropyranyloxy, pyrimidinyloxy, pyridyloxy, trifluorométhylpyridyloxy, benzothiazolyloxy, quinoléxyloxy, benzoyloxy(C₁ à C₈), benzènesulfonyloxy ou alkyl(C₁ à C₈)benzènesulfonyloxy.


3. Composé suivant la revendication 2, caractérisé en ce que Z représente un atome d'oxygène, R¹ représente un atome d'hydrogène, R² représente un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆ et R⁴, R⁵, X₁, X₂ et X₃ possèdent les significations qui leur ont été attribuées dans la revendication 2.
4. Composé suivant la revendication 1, caractérisé en ce que Z représente un atome d'oxygène, R¹ représente un atome d'hydrogène, R² représente un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆, R⁴ représente un atome d'hydrogène, R⁵ représente un atome d'hydrogène et A représente le radical phényle.
5. Composé suivant la revendication 1, caractérisé en ce que Z représente O, R¹ représente un atome d'hydrogène, R² représente le radical méthyle, R³ représente le radical méthyle, R⁴ représente un atome d'hydrogène, R⁵ représente un atome d'hydrogène et A représente le radical phényle.
6. Composé suivant la revendication 1, caractérisé en ce qu'il répond à la formule suivante :



dans laquelle R¹ et R² représentent chacun un atome d'hydrogène ou un radical alkyle en C₁ à C₆, R³ représente


un radical alkyle en C₁ à C₆, R⁴ et R⁵ représentent chacun un atome d'hydrogène, un radical alkyle en C₁ à C₆, alcoxy en C₁ à C₆, alkyle en C₁ à C₆ à substitution halogénée, silyle à substitution alkylique en C₁ à C₆, un atome d'halogène ou le radical nitro, R⁷ représente le radical phényle, un radical alcoxy(C₁ à C₆)phényle, halophényle, pyridyle ou pyrimidinyle et Z' représente -O-, -S-, -SO- ou NR- (R représentant un radical alkyle en C₁ à C₆).

7. Composé suivant la revendication 1, caractérisé en ce qu'il répond à la formule :




dans laquelle R¹ et R² représentent chacun un atome d'hydrogène ou un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆, A' représente le radical pyridyle ou pyrimidinyle, Z représente -O-, -OCH₂- ou -CH₂- et X₁ représente un atome d'hydrogène ou le radical trifluorométhyle.

8. Composé suivant la revendication 1, caractérisé en ce qu'il répond à la formule suivante :




dans laquelle R¹ et R² représentent chacun un atome d'hydrogène ou un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆ et Z représente -CH₂CH₂-,




-CH=CH-, -CH(OH)- ou -CO-.

9. Composé suivant la revendication 1, caractérisé en ce qu'il répond à la formule suivante :



15 dans laquelle R¹ et R² représentent chacun un atome d'hydrogène ou un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆ et A représente un radical alcényle comportant de 3 à 10 atomes de carbone, éventuellement substitué par pas plus de 3 atomes d'halogènes, ou un radical alcadiényle comportant de 3 à 10 atomes de carbone, éventuellement substitué par pas plus de 3 atomes d'halogènes.

20 10. Procédé de préparation d'un composé suivant l'une quelconque des revendications 1 à 9, caractérisé en, ce que l'on soumet un composé de la formule :



35 dans laquelle R⁴, R⁵, A et Z possèdent chacun les significations qui leur ont été précédemment attribuées dans la revendication 1, à une amidation ou une alcoxylation, ou vice versa.

40 11. Procédé suivant la revendication 10, caractérisé en ce que l'on entreprend d'abord l'amidation et ensuite l'alcoxylation.

12. Procédé suivant la revendication 10, caractérisé en ce que l'on entreprend d'abord l'alcoxylation et ensuite l'amidation.


45 13. Procédé suivant l'une quelconque des revendications 10 à 12, caractérisé en ce que l'on isole le composé ainsi obtenu et on le mélange à un ou plusieurs composants choisis parmi des véhicules ou supports, des diluants, de surfactifs, des agents d'adhérence ou d'adhésivité, des agents dispersifs et des stabilisants.

50 14. Composition ou agent fongicide, caractérisé en ce qu'il comprend, à titre d'ingrédient actif, au moins un composé suivant l'une quelconque des revendications 1 à 9, ainsi qu'un ou plusieurs composants choisis parmi les véhicules ou supports, les diluants, les surfactifs, les agents d'adhérence ou d'adhésivité, les agents dispersifs et les stabilisants, lesquels composants peuvent être inertes, la composition ou l'agent fongicide comprenant éventuellement aussi au moins un autre fongicide, un insecticide, un herbicide, ou un agent fertilisant ou engrais.

55 15. Procédé pour combattre des champignons phytopathogènes, caractérisé en ce que l'on applique, à titre d'ingrédient actif, au moins l'un des composés suivant l'une quelconque des revendications 1 à 9 à un endroit où des champignons phytopathogènes sont susceptibles de se propager ou se propagent.

16. Utilisation d'un composé suivant l'une quelconque des revendications 1 à 9 à titre de fongicide.

17. Utilisation d'un composé répondant à la formule suivante :



20 dans laquelle R^1 et R^2 représentent chacun indépendamment un atome d'hydrogène, un radical alkyle en C_1 à C_8 , ou cycloalkyle en C_3 à C_8 , R^3 représente un radical alkyle en C_1 à C_8 ou cycloalkyle en C_3 à C_8 , et R^4 et R^5 représentent chacun un atome d'hydrogène, un radical alkyle en C_1 à C_8 , alcoxy en C_1 à C_8 , alkyle en C_1 à C_8 à substitution halogénée, silyle à substitution alkylque en C_1 à C_8 , un atome d'halogène ou le radical nitro, A représente un radical pyridyle éventuellement substitué par pas plus de trois substituants et Z représente -O-, à titre de fongicide.


25 18. Composé suivant la revendication 3, caractérisé en ce que R^2 représente le radical méthyle, R^3 représente le radical méthyle, X_1 représente un atome d'hydrogène, X_2 représente un atome d'hydrogène et X_3 représente un atome d'hydrogène ou le radical 4-méthyle.

30 19. Composé suivant la revendication 2, caractérisé en ce que Z représente $-OCH_2-$, R^1 représente un atome d'hydrogène et R^2 représente le radical méthyle, R^3 représente le radical méthyle, R^4 et R^5 représentent chacun un atome d'hydrogène et X_1 représente un atome d'hydrogène et X_2 et X_3 représentent chacun un atome d'hydrogène, le radical méthyle, du chlore, ou le radical fluorométhyle.

35 20. Composé suivant la revendication 19, caractérisé en ce que X_2 et X_3 représentent chacun un atome d'hydrogène ou le radical méthyle.

40 21. Composé suivant la revendication 19, caractérisé en ce que X_2 représente un atome d'hydrogène et X_3 représente le radical 3- CF_3 , ou bien X_2 représente un atome d'hydrogène et X_3 représente le radical 2-Cl, ou bien X_2 représente un atome d'hydrogène et X_3 représente le radical 4-Cl.


22. Composé répondant à la formule suivante :



55 dans laquelle X représente O ou N-OH et A, Z, R^1 , R^2 , R^3 , R^4 et R^5 possèdent les significations qui leur ont été attribuées dans la revendication 1.

Revendications pour les Etats contractants suivants : ES, GR

1. Procédé de préparation d'un composé de la formule :



20 dans laquelle R^1 et R^2 représentent chacun indépendamment un atome d'hydrogène, un radical alkyle en C_1 à C_8 ou cycloalkyle en C_3 à C_8 , R^3 représente un radical alkyle en C_1 à C_8 ou cycloalkyle en C_3 à C_8 , R^4 et R^5 repré-

25 sentent chacun un atome d'hydrogène, un radical alkyle en C_1 à C_8 , un radical alcoxy en C_1 à C_8 , un radical alkyle en C_1 à C_8 à substitution halogénée, un radical silyle à substitution alkylque en C_1 à C_8 , un atome d'halogène, ou le radical nitro, A représente un groupe hydrocarboné insaturé, un radical hydrocarboné insaturé à substitution

halogénée, un radical phényle, ou un radical hétérocyclique, le radical phényle précité ou le radical hétérocyclique précité étant éventuellement substitué par pas plus de trois substituants et Z représente un radical $-CH_2-$, $-CH(OH)-$, $-CO-$, $-O-$, $-S-$, $-SO-$, $-NR-$ (R représentant un atome d'hydrogène ou un radical alkyle en C_1 - C_8), $-CH_2CH_2-$, $-CH=CH-$,




35 $-CH_2O-$, $-CH_2S-$, $-CH_2SO-$, $-OCH_2-$, $-SCH_2-$ ou $-SOCH_2-$, à condition que lorsque Z est O, A diffère du radical 2-pyridyle, caractérisé en ce que l'on soumet un composé de la formule :



50 dans laquelle R^4 , R^5 , A et Z possèdent chacun les significations qui leur ont été précédemment attribuées dans la revendication 1, à une amidation ou une alcoximation, ou vice versa.

- 55
2. Procédé suivant la revendication 1, caractérisé en ce que l'on entreprend d'abord l'amidation et ensuite l'alcoximation.
 3. Procédé suivant la revendication 1, caractérisé en ce que l'on entreprend d'abord l'alcoximation et ensuite l'amidation.

4. Procédé suivant l'une quelconque des revendications 1 à 3, caractérisé en ce que l'on choisit les réactifs pour obtenir un composé de la formule suivante :



dans laquelle Z représente -O-, -CH₂O-, -OCH₂-, -S-, -SO-, -SCH₂- ou -SOCH₂-, R¹ et R² représentent chacun un atome d'hydrogène ou un radical alkyle en C₁ à C₈, R³ représente un radical alkyle en C₁ à C₈, R⁴ et R⁵ représentent chacun indépendamment un atome d'hydrogène, un radical alkyle en C₁ à C₈, alcoxy en C₁ à C₈, un radical alkyle en C₁ à C₈ halogénosubstitué, un radical silyle à substitution alkylque en C₁ à C₈, un atome d'halogène, ou le radical nitro et X₁, X₂ et X₃ représentent chacun un atome d'hydrogène, un radical alkyle en C₁ à C₈, alcoxy en C₁ à C₈, alcanoyloxy en C₁ à C₈, alkyle en C₁ à C₈ à substitution halogénée, silyle à substitution alkylque en C₁ à C₈, un atome d'halogène, le radical nitro, cyano, un radical dialkyl(C₁ à C₈)phényle, phénylalcényloxy(C₂ à C₈), furylalcényloxy(C₂ à C₈), hydroxyle, alcynyloxy en C₂ à C₈, alcanoyloxy en C₁ à C₈, benzoyloxy, phénoxy, alcoxy(C₁ à C₈)phénoxy, nitrophénoxy, benzyloxy, cyanobenzyloxy, tétrahydropyranyloxy, pyrimidinyloxy, pyridyloxy, trifluorométhylpyridyloxy, benzothiazolyloxy, quinoléyloxy, benzoyloxy(C₁ à C₈), benzènesulfonyloxy ou alkyl(C₁ à C₈)benzènesulfonyloxy.

5. Procédé suivant la revendication 4, caractérisé en ce que Z représente un atome d'oxygène, R¹ représente un atome d'hydrogène, R² représente un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆ et R⁴, R⁵, X₁, X₂ et X₃ possèdent chacun les significations qui leur ont été attribuées dans la revendication 2.


6. Procédé suivant la revendication 1,

caractérisé en ce que Z représente un atome d'oxygène, R¹ représente un atome d'hydrogène, R² représente un radical alkyle en C₁ à C₆, R³ représente un radical alkyle en C₁ à C₆, R⁴ représente un atome d'hydrogène, R⁵ représente un atome d'hydrogène et A représente le radical phényle.

7. Procédé suivant la revendication 1,


Z représente un atome d'oxygène, R¹ représente un atome d'hydrogène, R² représente le radical méthyle, R³ représente le radical méthyle, R⁴ représente un atome d'hydrogène, R⁵ représente un atome d'hydrogène et A représente le radical phényle.

8. Procédé suivant la revendication 1, caractérisé en ce que l'on choisit les réactifs pour obtenir un composé de la formule suivante :




20 dans laquelle R^1 et R^2 représentent chacun un atome d'hydrogène ou un radical alkyle en C_1 à C_6 , R^3 représente un radical alkyle en C_1 à C_6 , R^4 et R^5 représentent chacun un atome d'hydrogène, un radical alkyle en C_1 à C_6 , alcoxy en C_1 à C_6 , alkyle en C_1 à C_6 à substitution halogénée, silyle à substitution alkylique en C_1 à C_6 , un atome d'halogène ou le radical nitro, R^7 représente le radical phényle, un radical alcoxy(C_1 à C_6)phényle, halophényle, pyridyle ou pyrimidinyle et Z' représente $-O-$, $-S-$, $-SO-$ ou $NR-$ (R représentant un radical alkyle en C_1 à C_6).

- 25 9. Procédé suivant la revendication 1, caractérisé en ce que l'on choisit les réactifs pour obtenir un composé de la formule suivante :




40 dans laquelle R^1 et R^2 représentent chacun un atome d'hydrogène ou un radical alkyle en C_1 à C_6 , R^3 représente un radical alkyle en C_1 à C_6 , A' représente le radical pyridyle ou pyrimidinyle, Z représente $-O-$, $-OCH_2-$ ou $-CH_2-$ et X_1 représente un atome d'hydrogène ou le radical trifluorométhyle.

- 45 10. Procédé suivant la revendication 1, caractérisé en ce que l'on choisit les réactifs pour obtenir un composé de la formule suivante :




dans laquelle R^1 et R^2 représentent chacun un atome d'hydrogène ou un radical alkyle en C_1 à C_6 , R^3 représente un radical alkyle en C_1 à C_6 et Z représente $-CH_2CH_2-$.



-CH=CH-, -CH(OH)- ou -CO-.

11. Procédé suivant la revendication 1, caractérisé en ce que l'on choisit les réactifs pour obtenir un composé de la formule suivante



dans laquelle R^1 et R^2 représentent chacun un atome d'hydrogène ou un radical alkyle en C_1 à C_6 , R^3 représente un radical alkyle en C_1 à C_6 et A représente un radical alcényle comportant de 3 à 10 atomes de carbone, éventuellement substitué par pas plus de 3 atomes d'halogène, ou un radical alcadiényle comportant de 3 à 10 atomes de carbone, éventuellement substitué par pas plus de 3 atomes d'halogènes.

12. Procédé suivant l'une quelconque des revendications précédentes, caractérisé en ce que l'on isole le composé résultant et on le mélange ensuite à un ou plusieurs composants choisis parmi les véhicules ou supports, les diluants, les surfactifs, les agents d'adhérence ou d'adhésivité, les agents dispersifs et les stabilisants.
13. Procédé de fabrication d'une composition fongicide, caractérisé en ce que l'on mélange, à titre d'ingrédient actif, au moins un composé tel que défini dans l'une quelconque des revendications 1 à 11, à un ou plusieurs composants choisis parmi les véhicules ou supports, les diluants, les surfactifs, les agents d'adhérence ou d'adhésivité, les agents dispersifs ou les stabilisants, lesquels composants pouvant être inertes, lesquels composants peuvent éventuellement comprendre en outre au moins un fongicide, insecticide, herbicide, ou fertilisant.
14. Procédé pour combattre des champignons phytopathogènes, caractérisé en ce que l'on produit un ingrédient actif par la mise en œuvre d'un procédé suivant l'une quelconque des revendications 1 à 12, et on applique ensuite l'ingrédient actif à un endroit où des champignons phytopathogènes se propagent ou sont susceptibles de se propager.